

**State: ANDHRA PRADESH**  
**Agriculture Contingency Plan for District: KRISHNA**

<b>1.0 District Agriculture profile</b>					
<b>1.1</b>	<b>Agro-Climatic/Ecological Zone</b>				
	Agro Ecological Sub Region (ICAR)	Eastern Coastal plane , hot, sub-humid to semi arid eco region (7.3,18.3)			
	Agro-Climatic Region (Planning Commission)	East Coast plain and hill region (XI)			
	Agro Climatic Zone (NARP)	Krishna Zone, RARS, Lam (AP-1)			
	List all the districts or part thereof falling under the NARP Zone	Guntur, Krishna and Prakasam			
	Geographic coordinates of district	Latitude	Longitude		Altitude
		15 <sup>0</sup> - 43 N and 17 <sup>0</sup> 10 N	80 E longitude and 81 <sup>0</sup> 33E		
	Name and address of the concerned ZRS/ ZARS/RARS/RRS/RRTTS	Regional Agricultural Research Station, Lam , Guntur, 522 034			
	Mention the KVK located in the district	Krishi Vigyan Kendra, Garikapadu, Krishna District 521 175			
<b>1.2</b>	<b>Rainfall</b>	Normal RF(mm)	Normal Rainy days (no)	Normal Onset ( specify week and month)	Normal Cessation (specify week and month)
	SW monsoon (June-Sep):	685	30-37	Second week of June	1 <sup>st</sup> week of October
	NE Monsoon (Oct-Dec):	250	7-10	2 <sup>nd</sup> week of October	Last week of December
	Winter (Jan- Feb)	16	0-5	-	-
	Summer (Mar-May)	83	2-4	-	-
	Annual	1034	41-60	-	-

<b>1.3</b>	<b>Land use pattern of the district</b> (latest statistics) 2009-10	Geographical area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	<b>Area ('000 ha)</b>	872.7	76.18	178.3	10.7	27.5	9.6	38.0	28.5	27.1

<b>1.4</b>	<b>Major Soils (common names like shallow red soils etc.,)</b>	<b>Area ('000 ha)</b>	<b>Percent (%) of total</b>
		1. Black Cotton Soils	260
	2. Red loams	76	18.76
	3.Sandy clay loams	41	10.12
	4.Alluvial Soils	11	2.74
	5. Saline soils	14	3.45
	Others (specify): Alkali	3	0.74

<b>1.5</b>	<b>Agricultural land use</b>	<b>Area ('000 ha)</b>	<b>Cropping intensity %</b>
	Net sown area	463.0	154.7
	Area sown more than once	253.3	
	Gross cropped area	716.3	

<b>1.6</b>	<b>Irrigation</b>	Area ('000 ha)		
	Net irrigated area	340		
	Gross irrigated area	520		
	Rainfed area	170		
	<b>Sources of Irrigation</b>	Number	Area ('000 ha)	Percentage of total irrigated area
	Canals	KE canal System	251	73.82
	Tanks	983	38	11.18
	Open wells	15552		
	Bore wells	13549	30	8.82
	Lift irrigation			
	Micro-irrigation			
	Other sources	64	21	6.18
	Total Irrigated Area		340	100.0
	Pump sets			
	No. of Tractors	8469		
	<b>Groundwater availability and use* (Data source: State/Central Ground water Department /Board)</b>	No. of blocks/ Tehsils	(% ) area	
	Over exploited		not available	
	Critical			
	Semi- critical			
	Safe			
Wastewater availability and use				
Ground water quality	not available			
*over-exploited: ground water utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%				

**Area under major field crops & horticulture etc. (2009-10)**

1.7	Major Field Crops cultivated	Area ('000 ha)					
		Kharif		Rabi		Summer	Total
		Irrigated	Rainfed	Irrigated	Rainfed		
1	Paddy	228.74		39.98	-		268.72
2	Blackgram	-	2.51	107.11	24.69	-	134.31
3	Maize	2.6	1.8	20.35	-		24.75
4	Cotton	48.23	-	-	-	-	48.23
5	Greengram	-	4.03	-	7.62	-	11.65
6	Sugarcane	-	-	13.13	-	-	13.3
7	Chillies	6.53		0.72	-	-	11.56
8	Groundnut		1.03	1.0		-	2.03
9	Tobacco	-	-	1.59		-	1.59
10	Red gram	-	1.62	-	-	-	1.62
	<b>Horticulture crops - Fruits</b>	<b>Total area ('000 ha)</b>					
1	Mango	63.5					
2	Banana	2.7					
3	Guava	1.2					
4	Papaya	0.1					
5	Batavia	0.04					
6	Cashew nut	0.3					
	<b>Horticultural crops - Vegetables</b>	<b>Total area('000 ha)</b>					
1	Chillies	7.9					
2	Bhendi	2.0					
3	Tomato	1.6					
4	Gourds	1.1					
5	Cucumber	1.1					

6	Leaf Vegetables	0.3
---	-----------------	-----

	<b>Medicinal and Aromatic crops</b>	<b>Total area('000 ha)</b>
1	Turmeric	1.8
2	Ginger	0.02
3	Onion	0.02
4	Eucalyptus	0.9
5	Betel vine	0.2
	<b>Plantation crops</b>	<b>Total area('000 ha)</b>
1	Banana	1.4
2	Coconut	2.1
3	Sapota	0.4
4	Acid lime	0.7
5	Oil palm	3.1
	<b>Fodder crops</b>	<b>Total area('000 ha)</b>
1	Jowar	1.7
2	Maize	1.1
3	Para napier	1.3
4	Pillipaesara	6.2
5	Sunhemp	
	<b>Total fodder crop area</b>	10.2
	<b>Grazing land</b>	
	<b>Sericulture etc</b>	
	<b>Others (Specify)</b>	

<b>1.8</b>	<b>Livestock</b>		<b>Male (number)</b>	<b>Female (number)</b>	<b>Total (number)</b>		
	Non descriptive Cattle (local low yielding)		42.9	50.5	93.5		
	Crossbred cattle		1.9	9.0	10.9		
	Non descriptive Buffaloes (local low yielding)		117.6	801.8	919.4		
	Graded Buffaloes						
	Goat				156.0		
	Sheep				482.1		
	Others (Camel, Pig, Yak etc.)				16.77		
	Commercial dairy farms (Number)						
<b>1.9</b>	<b>Poultry</b>		<b>No. of farms</b>	<b>Total No. of birds (number)</b>			
	Commercial						
	Backyard						
<b>1.10</b>	<b>Fisheries (Data source: Chief Planning Officer)</b>						
	<b>A. Capture</b>						
	i) Marine (Data Source: Fisheries Department)	No. of fishermen	Boats		Nets		Storage facilities (Ice plants etc.)
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Gill nets)	Non-mechanized (Shore Seines, Stake & trap nets)	
		5774	96	781 / 137	214 / 72375	366 / 10	32 / 0
	ii) Inland (Data Source: Fisheries Department)	No. Farmer owned ponds		No. of Reservoirs		No. of village tanks	
		9131		1		219	
	<b>B. Culture</b>						
			<b>Water Spread Area (ha)</b>	<b>Yield (t/ha)</b>	<b>Production ('000 tons)</b>		
	i) Brackish water (Data Source: MPEDA/ Fisheries Department)		4750	-	6.4		
	ii) Fresh water (Data Source: Fisheries Department)		23958	-	10.2		
Others			-	394.3			



1.12	Sowing window for 5 major field crops (start and end of normal sowing period)	Paddy	Maize	Pulses	Groundnut	Cotton
	<i>Kharif</i> - Rainfed	-	July 1 <sup>st</sup> fortnight – July 2 <sup>nd</sup> fortnight	June 1 <sup>st</sup> fortnight – July 2 <sup>nd</sup> fortnight	-	June 1 <sup>st</sup> fortnight – July 2 <sup>nd</sup> fortnight
	<i>Kharif</i> -Irrigated	June 1 <sup>st</sup> fortnight – July 2 <sup>nd</sup> fortnight	July 1 <sup>st</sup> fortnight – July 2 <sup>nd</sup> fortnight	-	-	-
	<i>Rabi</i> - Rainfed	-	-	September 1 <sup>st</sup> fortnight – October 1 <sup>st</sup> fortnight	October 2 <sup>nd</sup> fortnight – November 1 <sup>st</sup> fortnight	-
	<i>Rabi</i> -Irrigated	December 2 <sup>nd</sup> fortnight – January 1 <sup>st</sup> fortnight	-	October 2 <sup>nd</sup> fortnight	November 2 <sup>nd</sup> fortnight – December 1 <sup>st</sup> fortnight	-

1.13	What is the major contingency the district is prone to? (Tick mark and mention years if known during the last 10 year period)	Regular	Occasional	None
	Drought		√	
	Flood		√	
	Cyclone	√		
	Hail storm			
	Heat wave			
	Cold wave			
	Frost			
	Sea water intrusion		√	
	Pests and diseases (specify)	<u>Rice</u> : Blast, planthoppers, stem rot, leaf folder, sheath blight <u>Redgram</u> : Maruca and Helicoverpa, sterility mosaic, wilt, pod fly <u>Cotton</u> : Sucking pest complex, <u>pink</u>		



		<u>boll worm, root rots, wilt</u> Blackgram : YMV, leaf crinkle virus, stem canker, Maruca pod borer, Spodoptera, powdery mildew Maize: fall army worm, Late wilt, stem borers Sugar cane: early shoot borer, whip smut, scales, root grub Groundnut: leaf spots, stem rot, root rot, foot rot, thrips		
	Others (Fog)			

<b>1.14</b>	<b>Include Digital maps of the district for</b>	Location map of district within State as Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure 2	Enclosed: Yes
		Fertility Status as Annexure 3	Enclosed: Yes

### KRISHNA DISTRICT INDEX MAP






Fig: Location map of the district



### SOIL MAP - KRISHNA DISTRICT



## 2.0 Strategies for weather related contingencies

### 2.1 Drought

#### 2.1.1 Rainfed situation

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 2 weeks (3 <sup>rd</sup> week of June)	Black soils – Rainfed	Cotton	No change	Normal practices	-
		Greengram			
		Redgram			
	Red soils – Rainfed	Cotton			
		Redgram (Sole crop)			
		Redgram+Greengram / Groundnut (1:7)			
		Green gram/ black gram – Red gram (7:1)			
Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 4 weeks (July 1 <sup>st</sup> week)	Black soils – Rainfed	Cotton	No change	Normal practices	
	Red soils – Rainfed	Cotton		Normal practices	
		Redgram (Sole crop)		Reduce Redgram row spacing 180 cm to 150 cm	
		Redgram+Greengram / Groundnut (1:7)		Normal practices	
		Green gram/ black gram – Red gram (7:1)			

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 6 weeks (July 3 <sup>rd</sup> week)	Black soils – Rainfed	Cotton	No change	Adopt closer spacing of 90x45cms	-
		Redgram	Short duration varieties PRG-100, ICPL-84031		
	Red soils - Rainfed	Cotton	No change	Adopt closer spacing of 90X45 cm	
		Redgram (Sole crop)		Reduce row spacing 180 cm to 150 cm	
		Redgram+Greengram/ Groundnut		Normal practices	

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 8 weeks (August 1 <sup>st</sup> week)	Black soils – Rainfed	Cotton	No change	Adopt closer spacing of 90X 45 cm	-
	Red soils - Rainfed	Cotton		Adopt closer spacing of 90 X 30 cm. Top dressing of fertilizer at 20 days interval	
		Redgram (Sole crop)		Reduce row spacing 180 cm to 120 cm	
		Redgram+Greengram/ Groundnut	Redgram sole crop	Reduce row spacing 180 cm to 120 cm	

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Early season drought (Normal onset)					
Normal onset followed by 15-20 days dry spell after sowing leading to poor germination/crop stand etc.	Black soils – Rainfed	Cotton	Gap filling to be done by pot watering 7- 10 days after sowing if the crop stand is poor	When the crop is 2 weeks old, take up intercultivation to conserve moisture Spray 2 % urea solution or 1 % water soluble fertilizers like 19-19-19, 1% KNO <sub>3</sub> to supplement nutrition	-
	Red soils - Rainfed	Cotton			
		Redgram (sole crop)		Making of deep conservation furrow or Inter cultivation to be done after 2 weeks of sowing or mulching to conserve soil moisture Foliar spray of 2% urea to supplement nutrition	
		Redgram+ Greengram	-	Foliar spray of 2% urea to supplement nutrition	

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)					
At vegetative stage	Black soils – Rainfed	Cotton	Spray 2 % urea solution or 1 % water soluble fertilizers like 19-19-19 or 1 % KNO <sub>3</sub>	Inter cultivate periodically (7-10 days interval) or mulching to conserve soil moisture	-
	Red soils - Rainfed	Cotton		Inter cultivate	

				periodically (7-10 days interval) or mulching to conserve soil moisture	
		Redgram (sole crop)		Making of deep conservation furrow or Inter cultivation to be done after 2 weeks of sowing Making of deep conservation furrow or to conserve soil moisture Foliar spray of 2% urea to supplement nutrition	
		Redgram+Greengram	Harvest intercrops as fodders as chances of grain yield are poor Supplement the nutrients to the main crop through foliar spray	--	

Condition			Suggested Contingency measures		
Mid season drought (long dry spell)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
At reproductive stage	Black soils – Rainfed	Cotton	Spray urea - 2 % or KNO <sub>3</sub> 1% or other water soluble fertilizers like 19-19-19 to supplement nutrition	Intercultivation to create soil mulch to conserve moisture	-
	Red soils - Rainfed	Cotton		Intercultivation to conserve moisture. Supplemental irrigation, if available	
		Redgram (sole crop)			
		Redgram+Greengram		--	



Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Rabi Crop planning	Remarks on Implementation
Terminal drought	Black soils rainfed	Cotton	Spray urea - 2 % or KNO <sub>3</sub> 1% or other water soluble fertilizers 1 % to supplement nutrition Topping to prevent formation of new vegetative and reproductive flush Supplemental irrigation if available	Blackgram/Cowpea/Castor/Minor millets/Bengalgram	-
	Red soils - Rainfed	Cotton		Cowpea/Castor/Minor millets	
		Redgram (sole crop)			
	Redgram+Greengram				

### 2.1.2 Irrigated situation

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delayed release of water in canals due to low rainfall	Black soils – Canal irrigated (KED)	Green manure – Rice – Blackgram/Maize	Green manure/short duration greengram – Rice – Greengram / Blackgram / Maize	Growing of short duration greengram like GGG-1, LGG-460,WGG-42 Increase the plant population. Adopt prophylactic measures for Blast During Rabi season select short duration Blackgram varieties like LBG 20, LBG 752, GBG-1,PU-31,TBG-104 Greengram can be grown in rice fallows under late seasonal conditions instead of blackgram Zero tillage maize in paddy fields with varieties DHM 117, Trishulatha and other popular hybrids	

Condition	Suggested Contingency measures				
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
		Paddy – Sugarcane (plant) and Sugarcane (ratoon)	Paddy – Sugarcane plant and ratoon – paddy	Raising of nurseries with single buded sets to save the time and water Use of drip system to save the water quantity Mulching with sugarcane trash between rows and frequent intercultivations to conserve moisture	
	Red Soils/Black Soils – Canal irrigated (NSP left canal Command area/ Tank fed and lift irrigation )	Greengram – Rice – Greengram / Maize / Blackgram / Fodder	Rice- Greengram/ Maize/ Blackgram/ Jowar	Management of aged seedlings Direct sowing of short duration varieties viz., MTU 1153, MTU 1156, MTU 1121, NLR 34449 etc. Short duration crops like greengram, blackgram, maize and groundnut	

Condition	Suggested Contingency measures				
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Limited release of water in canals due to low rainfall	Black soils – Canal irrigated (KED)	Green manure – Rice – Blackgram/Maize	Green manure – Rice – Black gram/Greengram - Aerobic rice	Management of over aged seedlings Direct seeding varieties (<135 days) with short duration Adopt alternate wetting and drying upto primordial Initiation stage to save water Short duration varieties of crops shall be selected. Blackgram varieties LBG 20, LBG 752, GBG-1,PU-31,TBG-104 and maize varieties DHM 117, Trishulatha Water saving micro irrigation systems	
		Paddy – Sugarcane plant and ratoon – paddy	No change	Raising of nurseries with single budded sets to save the time and water Conservation practices like inter cultivation, earthing up, Alternate row irrigation shall be practiced Water loss during conveyance can be reduced by using PVC/Metallic pipes instead of running water in open field channels	
	Black soils/Red soils – Left Canal irrigated (NSP left canal Command area/ Tank fed and lift irrigation )	Greengram – Rice – Blackgram/ Greengram/Maize/ Fodder	<ul style="list-style-type: none"> <li>• Green manure – Rice – Greengram/ Blackgram/Jowar / Fodder</li> <li>• Redgram + Greengram /Jowar</li> <li>• Cotton (Wherever drainage facilities available)</li> </ul>	For rice and rice fallow crops the agronomic measures as suggested for the above farming situation shall be followed  Proper drainage facilities should be created to take up cropping systems as suggested	

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Non release of water in canals under delayed onset of monsoon in catchment	Black soils – Canal irrigated (KED)	Green manure – Rice – Blackgram/Maize	<ul style="list-style-type: none"> <li>• Go for early rabi</li> <li>• Green manure – Blackgram – Groundnut/Jowar</li> </ul>	<p>Green manure crops should be incorporated</p> <p>Sowing of ID crops (blackgram and groundnut) can be taken from September second fortnight onwards</p> <p>Blackgram can be grown from December to February with two to three irrigations after the harvest of early Rabi crops</p>	Farmers are to be trained on the upland crop cultivation practices
	Black soils/Red soils – Canal irrigation (NSP left canal Command area/ Tank fed and lift irrigation )	<ul style="list-style-type: none"> <li>• Greengram/Green manure – Rice – Blackgram/Greengram/Jowar/Fodder</li> </ul>	<ul style="list-style-type: none"> <li>• Greengram/Green manure – Blackgram / Jowar</li> <li>• Green manure/ Greengram – Cotton</li> </ul>	<p>Green manure crops should be incorporated</p> <p>Sowing of upland crops can be taken from September second fortnight onwards</p> <p>Blackgram can be grown from December to February/March with two to three irrigations after the harvest of early rabi crops</p>	

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Lack of inflows into tanks due to insufficient /delayed onset of monsoon	Green manure/ fodder/Paddy	Green gram / fodder crops and green manure crops – medium to short duration paddy	Green manure crops - ID crops like maize , jowar , red gram, groundnut	<ul style="list-style-type: none"> <li>• Green manure crops should be incorporated in to the soil at right stage and allow it to decompose with the moisture received from rain</li> <li>• Sowing of crops can be taken from September second fortnight onwards</li> <li>• Maize, Blackgram can be grown from December to February with two to three irrigations after the harvest of early rabi crops</li> </ul>	

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures	Remarks on Implementation
Insufficient groundwater recharge due to low rainfall	NA				
Any other condition (specify)					

**2.2 Unusual rains (untimely, unseasonal etc)** (for both rainfed and irrigated situations)

<b>Condition - Continuous high rainfall in a short span leading to water logging and Heavy rainfall with high speed winds in a short span</b>				
<b>Crop</b>	<b>Suggested contingency measure</b>			
	<b>Vegetative stage</b>	<b>Flowering stage</b>	<b>Crop maturity stage</b>	<b>Post harvest</b>
Rice	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up gap filling either with available nursery or by splitting the tillers from the surviving hills</p> <p>Take up suitable plant protection measures in anticipation of pest &amp; disease out breaks like army worm, sheath blight etc.</p>	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up suitable plant protection Measures in anticipation of pest &amp; disease out breaks like leaf folder, sheath rot, false smut etc.</p>	<p>Drain the excess water as early as possible</p> <p>Take up suitable plant protection measures in anticipation of pest &amp; disease out breaks like climbing cut worm</p>	<p>Drain out water and spread sheaves loosely in field or field bunds where there is no water stagnation</p> <p>Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds</p> <p>Thresh after drying the sheaves properly</p> <p>Ensure proper grain moisture before storing</p>
Cotton	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up inter cultivation at optimum soil moisture condition to loosen and aerate the soil and to control weeds</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19to support nutrition</p> <p>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</p> <p>Take up timely control measures against the out break of pests like Spodoptera, Helicoverpa etc.</p>	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19at 1% to support nutrition</p> <p>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</p> <p>Take up timely control measures against the out break of pests like Spodoptera, Helicoverpa etc.</p>	<p>Drain the excess water as early as possible</p> <p>Pray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19at 1% to support nutrition</p> <p>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</p> <p>Take up timely control measures against the out break of pests like Spodoptera, Helicoverpa etc.</p>	<p>Dry the produce properly before packing and sending to market</p>
Redgram	Drain the excess water as early as	Drain the excess water as early as	Drain the excess water as early	Spread the bundles

	<p>possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up inter cultivation at optimum soil moisture condition to loosen and aerate the soil and to control weeds</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition</p>	<p>possible</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition</p> <p>Take up timely control measures against the out break of pests like Spodoptera, Helicoverpa etc.</p>	<p>as possible</p> <p>Allow the crop to dry completely before harvesting</p>	<p>drenched in rain on field bunds or drying floors to quicken the drying</p> <p>Thresh the bundles after they are dried properly</p> <p>Dry the grain to proper moisture per cent before bagging and storing to prevent deterioration in quality during storage</p>
Blackgram	<p>Drain the excess water as early as possible</p> <p>Apply 4-5 kg N /ha after draining excess water</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-at 1% to support nutrition. Correction of iron deficiency by spraying 0.5% FeSO<sub>4</sub></p> <p>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</p> <p>Take up timely control measures against the out break of pests like <i>Spodoptera</i> etc.</p>	<p>Drain the excess water as early as possible</p> <p>Apply 4-5 kg N /ha after draining excess water</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19at 1% to support nutrition. Correction of iron deficiency by spraying 0.5% FeSO<sub>4</sub></p> <p>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</p> <p>Take up timely control measures against the out break of pests like <i>Spodoptera</i> etc.</p>	<p>Drain the excess water as early as possible</p> <p>Allow the crop to dry completely before harvesting</p>	<p>Spread the bundles drenched in rain on field bunds or drying floors to quicken the drying</p> <p>Thresh the bundles after they are dried properly</p> <p>Dry the grain to proper moisture per cent before bagging and storing to prevent deterioration in quality during storage</p>
Maize	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water. Correction of iron deficiency by spraying 0.5% FeSO<sub>4</sub>+ Citric Acid 0.1 % solution 2-3 times</p> <p>Take up intercultivation and at optimum soil moisture condition to loosen and aerate the soil and to control weeds</p> <p>Earthenup the crop for anchorage</p>	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water. Correction of iron deficiency by spraying 0.5% FeSO<sub>4</sub>+ Citric Acid 0.1 % solution 2-3 times</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19at 1% to support nutrition</p> <p>Take up timely control measures for sheath blight and post flowering</p>	<p>Drain the excess water as early as possible</p> <p>Allow the crop to dry completely before harvesting</p>	<p>Harvest the cobs after the they are dried up properly.</p> <p>Dry the grain to optimum moisture condition before storing</p>

	<p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19at 1% to support nutrition</p> <p>Take up timely control measures for Pink stem borer, sheath blight and Turcicum leaf blight</p>	stalk rots		
<b>Horticulture crops – Fruits</b>				
Mango	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the mature produce in a clear sunny day'</p>	<p>Store the fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p>
Banana	<p>Drain the excess water as soon as possible</p> <p>Inter-cultivate the soil with gorru for aeration.</p> <p>Spray 0.5 % KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 80 g MOP + 100 g Urea per plant at two to three times intervals.</p> <p>Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.</p> <p>If the age of the plant is less than three months and submergence up to three feet better to replant the garden.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray 0.5 % KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 80 g MOP + 100 g Urea per plant at two to three times intervals.</p> <p>If the age the plant is more than three months and less than seven months allow one sword sucker for ratoon and take up fertilization at monthly intervals for four months.</p> <p>Staking with bamboos to prevent further lodging.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the marketable bunches in a clear sunny day.</p> <p>Spray 0.5 % KNO<sub>3</sub> or Urea 2% solution 2-3 times for quick development of immature bunches.</p> <p>Staking with bamboos to prevent further lodging.</p>	<p>Use ripening chambers for quick ripening</p> <p>Market the produce as soon as possible.</p>
Guava	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the mature produce as soon as possible.</p>	<p>Store the produce in well-ventilated place temporarily before it can be marketed.</p> <p>Market the produce as soon as possible.</p>
<b>Horticultural crops - Vegetables</b>				



Chillies	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.</p> <p>In case of severe damage (considered as complete economical loss), and the contingency period is between June to August, sowing of best alternative crop must be taken up.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the matured fruits in a clear sunny day.</p>	<p>Dry the pods on concrete floor immediately after the appearance of sunlight (or).</p> <p>Use poly house solar driers for quick drying</p> <p>Grade the pods and market as soon as possible.</p> <p>Do not store such produce for long periods.</p>
Tomato	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 12 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Spray COC 30 g in 10 liters of water, 2-3 times against leaf spots.</p> <p>Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.</p> <p>In case of severe damage (considered as complete economical loss), and the contingency period is between June to August, sowing of best alternative crop must be taken up.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 10 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Spray COC 30 g in 10 liters of water, 2-3 times against leaf spots</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the marketable fruits in a clear sunny day'</p>	<p>Store the harvested fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p>
Cucumber	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 10 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Gap filling may be taken up if the</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 10 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution once.</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature produce as soon as possible.</p> <p>Store the produce in well ventilated place temporarily before it can be marketed.</p>

	plants are two weeks old and sowing window is still available for the crop. In case of severe damage (considered as complete economical loss), and the contingency period is between June to August, sowing of best alternative crop must be taken up.			Market the produce as soon as possible.
Plantation crops				
Oil palm	Planting should be done on mounts or bunds Drainage system, suited to local conditions may be provided to remove surplus water from root zone Relief drains [shallow] channels are opened at places where water accumulates and connected with main drain to remove water from the surface	Drain the excess water as soon as possible Apply booster dose of NPK fertilizers	Drain the excess water as soon as possible Apply booster dose of NPK fertilizers Harvest the mature nuts as soon as possible	Store the produce in well ventilated place temporarily before it can be market Market the nuts as soon as possible.
Turmeric	Drain the excess water as soon as possible Spray Urea 2% or 1% KNO <sub>3</sub> followed by Ferrous Sulphate 0.5% + Citric Acid 0.1 % solution 2-3 times. In case of severe damage (considered as complete economical loss or if inundation is more than for four days), and the contingency period is between June to August, sowing of best alternative crop must be taken up.	Drain the excess water as soon as possible Spray Urea 2% or 1% KNO <sub>3</sub> solution 2-3 times.	Drain the excess water as soon as possible Harvest the rhizomes when field comes to normal	Dry the rhizomes on concrete floor or use boilers (if available) for processing immediately Grade and separate the rotten and mould affected rhizomes. Pack the dried material in gunny bags disinfected with safe insecticides Store in a well ventilated rooms
Coconut	Planting should be done on mounts or bunds Drainage system, suited to local conditions may be provided to remove surplus water from root zone Relief drains [shallow] channels are opened at places where water	Drain the excess water as soon as possible Apply booster dose of NPK fertilizers	Drain the excess water as soon as possible Apply booster dose of NPK fertilizers Harvest the mature nuts as soon as possible.	Store the produce in well ventilated place temporarily before it can be market Market the nuts as soon as possible.

	accumulates and connected with main drain to remove water from the surface			
<b>Condition - Outbreak of pests and diseases due to unseasonal rains</b>				
Rice	Stem rot and Sheath blight - need based plant protection measures to be initiated based on incidence levels	BPH, Blast, Sheath blight incidence may increase due to unseasonal rains - need based plant protection measures to be initiated	Climbing cutworm and neck blast	-
Cotton	Jassids, Wilt and root rot, Bacterial leaf blight - Need based plant protection measures to be initiated	Jassids, <i>Spodoptera</i> , Wilt and root rot, Bacterial leaf blight, Grey mildew - Need based plant protection measures to be initiated	Dusky cotton bug, Grey mildew - Need based plant protection measures to be initiated	Dry the seed cotton properly after picking and store it under shade in aerated place
Redgram	Wilt and root rot - Need based plant protection measures to be initiated	Wilt and root rot. Need based plant protection measures to be initiated	-	
Blackgram	Spodoptera - Need based plant protection measures to be initiated	Spodoptera, Leaf spots, Powdery mildew - Need based plant protection measures to be initiated	Spodoptera, Rust - Need based plant protection measures to be initiated	
Maize		Jassids, Wilt and Stalk rot	Post flowering Stalk rots may aggravate if unseasonal rains occurs	

### 2.3 Floods

<b>Condition</b>	<b>Transient water logging/ partial inundation and Continuous submergence for more than 2 days</b>			
	<b>Suggested contingency measure</b>			
	<b>Seedling / nursery stage</b>	<b>Vegetative stage</b>	<b>Reproductive stage</b>	<b>At harvest</b>
Rice	Drain out the excess water at the earliest Apply booster dose of 0.2 kg N/40 sq. m Takeup proper weed control measures Spray of ZnSO <sub>4</sub> , FeSO <sub>4</sub> to correct micronutrient	Drain out the excess water at the earliest Take up gap filling either with available nursery or by splitting the tillers from the surviving hills Apply a booster dose of 20 kg N/acre Spray ZnSO <sub>4</sub> 0.2% if it is less than 45 days after transplanting Takeup need based plant protection	Drain out the excess water at the earliest Takeup need based plant protection measures	Drain out water .Spread sheaves loosely in field or field bunds where there is no water stagnation Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds

	deficiencies	measures		Thresh after drying the sheaves properly Ensure proper grain moisture before storing
Cotton	<p>Drain out the excess water at the earliest.</p> <p>Take up the gap filling at the earliest.</p> <p>Inter cultivate at optimum field moisture condition.</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water.</p> <p>To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition.</p> <p>Take up plant protection measures against possible pests and disease incidence.</p> <p>Mortality is most likely hence resowing to be taken up.</p> <p>With short duration hybrids Adopt closer spacing of 90X45 or 90X30 cm.</p>	<p>Drain out the excess water at the earliest</p> <p>Inter cultivate at optimum field moisture condition.</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water.</p> <p>To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition.</p> <p>Spray of micronutrients two times at 7-10 days interval.</p> <p>Take up plant protection measures against possible pests and disease incidence.</p>	<p>Drain out the excess water at the earliest.</p> <p>To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition.</p> <p>Take up plant protection measures against possible pests and disease incidence.</p>	<p>Kapas picking should be done carefully to prevent admixtures with waste plant material.</p> <p>To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition.</p>
Sugarcane	<p>Drain out the excess water at the earliest</p>	<p>Drain out the excess water at the earliest</p> <p>Apply 50 kg N urea and 25k / ha</p>	<p>Drain out the excess water at the earliest</p> <p>Apply 50 kg N urea and 20k/ha</p>	<p>Drain out the excess water at the earliest</p> <p>Harvest the crop as early as possible</p>
Redgram	<p>Drain out the excess water at the earliest</p> <p>Take up the gap filling at the earliest</p> <p>Inter cultivate at optimum field moisture condition</p>	<p>Drain out the excess water at the earliest</p> <p>Take up the gap filling at the earliest</p> <p>Inter cultivate at optimum field moisture condition</p> <p>Apply 4-5 kg N/acre after draining excess water</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble</p>	<p>Drain out the excess water at the earliest</p> <p>To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>Drain out the excess water at the earliest</p> <p>Harvest the crop when the field condition permits</p> <p>Drying of bundles should be done on elevated places like filed bunds or drying floors</p>

	Apply 4-5 kg N/acre after draining excess water	fertilizers like 19-19-19 at 1% to support nutrition Proper weed control measures to be taken up Need based plant protection measures to be taken up	Apply 20 kg N + 10 kg K /acre after draining excess water Need based plant protection measures to be taken up	
Blackgram	Drain out the excess water at the earliest Take up the gap filling at the earliest Take up weed control either mechanically or through weedicides Apply 4-5 kg N/acre after draining excess water Take up plant protection measures against possible pests and disease incidence	Drain out the excess water at the earliest Take up weed control either mechanically or through weedicides Apply 4-5 kg N/acre after draining excess water To spray KNO <sub>3</sub> 1 % or water soluble fertilizers like 19-19-19at 1% to support nutrition Take up plant protection measures against possible pests and disease incidence	Drain out the excess water at the earliest Apply 4-5 kg N/acre after draining excess water To spray KNO <sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition Take up plant protection measures against possible pests and disease	Drain out the excess water at the earliest Harvest the crop after the fields are dried up  Dry the bundles on field bunds and drying floors Dry the grain to optimum moisture content before storage
Maize	Drain out the excess water at the earliest Take up weed control either mechanically or through weedicides Intercultivation and earthing up to be done Apply 20 kg N + 10 kg K /ha after draining excess water Take up plant protection measures against possible pests and disease incidence  Re - sow the crop if mortality is > 15 %	Drain out the excess water at the earliest Take up weed control either mechanically or through weedicides Intercultivation and earthing up to be done Apply 20 kg N + 10 kg K /ha after draining excess water Take up plant protection measures against possible pests and disease incidence  Spray KNO <sub>3</sub> @ 1 % or water soluble fertilizers like 19-19-19 @ 1% to support nutrition Need based plant protection measures to be taken up	Drain out the excess water at the earliest Take up plant protection measures against possible pests and disease incidence  Apply 20 kg N + 10 kg K /acre after draining excess water Spray KNO <sub>3</sub> @ 1 % or water soluble fertilizers like 19-19-19 @ 1% to support nutrition Need based plant protection measures to be taken up	Drain out the excess water at the earliest Cob picking to be done after they are dried fully
<b>Horticulture</b>				
mango	Drain the excess water as soon	Drain the excess water as soon as	Drain the excess water as	Drain the excess water as soon

banana	as possible	possible	soon as possible	as possible.
guava	Spray 1% KNO <sub>3</sub> or Urea 2% solution 2-3 times.	Spray 1% KNO <sub>3</sub> or Urea 2% solution 2-3 times.	Spray 1% KNO <sub>3</sub> or Urea 2% solution 2-3 times.	Harvest the mature fruits as soon as possible. Store the fruits in well ventilated place temporarily before it can be marketed. Market the fruits as soon as possible. Spray Dithane M-45 0.3% or bavistin 0.1% against Anthracnose

#### 2.4 Extreme events: Heat wave / Cold wave / Frost / Hailstorm / Cyclone ✓

Extreme event type	Suggested contingency measure			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Heat Wave				
	Cyclones are common in this district, lot of damage occurred in all crops			
Cold wave				
Frost				
Hailstorm				
Cyclone				
Rice	Drain out the excess water at the earliest Apply booster dose of 0.2 kg N/40 sq. m Spray micronutrients like Zn, Fe 2-3 times at 4 -5 days interval Takeup proper weed control measures	Drain out the excess water at the earliest Apply booster dose of 20 kg N/Acre Spray ZnSO <sub>4</sub> 0.2 % if it is less than 45 days after transplanting Take up need based plant protection measures	Drain out the excess water at the earliest Takeup need based plant protection measures Lodged plants to be lifted and tied together to make them stand erect	Drain out water spread sheaves loosely in field or field bunds where there is no water stagnation Spray common salt at 5% to prevent germination of seed and spoilage of straw from moulds Thresh after drying the sheaves properly 4. Ensure proper grain moisture before storing
Cotton	Drain out the excess water at the	Drain out the excess water at the	Drain out the excess	Kapas picking should be done

	<p>earliest</p> <p>Inter cultivate at optimum field moisture condition</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p>	<p>earliest</p> <p>Inter cultivate at optimum field moisture condition</p> <p>Earthing up to be done to provide anchorage to plants</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>To spray KNO<sub>3</sub>@1 % or water soluble fertilizers like 19-19-19@ 1% to support nutrition</p> <p>Spray of micronutrients two times at 7-10 days interval</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>water at the earliest</p> <p>Spray KNO<sub>3</sub> @1 % or water soluble fertilizers like 19-19-19 @ 1% to support nutrition</p> <p>Earthing up to be done to provide anchorage to plants</p> <p>Spray of micronutrients two times at 7-10 days interval</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>carefully to prevent admixtures with waste plant material</p>
Redgram	<p>Drain out the excess water at the earliest</p> <p>Inter cultivate at optimum field moisture condition</p> <p>Apply 4-5 kg N/acre after draining excess water</p>	<p>Drain out the excess water at the earliest</p> <p>Inter cultivate at optimum field moisture condition</p> <p>Apply 4-5 kg N/acre after draining excess water</p>	<p>Drain out the excess water at the earliest</p> <p>To spray KNO<sub>3</sub>@ 1% or water soluble fertilizers like 19-19-19 @ 1% to support nutrition</p> <p>3. Take up plant protection measures against possible pests and disease incidence</p>	<p>Drain out the excess water at the earliest</p> <p>Harvest the crop when the field condition permits</p> <p>Drying of bundles should be done on elevated places like filed bunds or drying floors</p>
Blackgram	<p>Drain out the excess water at the earliest</p> <p>Takeup weed control either mechanically or through weedicides</p> <p>Apply 4-5 kg N/acre after draining excess water</p>	<p>Drain out the excess water at the earliest</p> <p>Take up weed control either mechanically or through weedicides</p> <p>Apply 4-5 kg N/acre after draining excess water</p> <p>Spray KNO<sub>3</sub> @1 % or water soluble fertilizers like 19-19-19 @ 1% to support nutrition</p> <p>Take up plant protection measures against possible pests</p>	<p>Drain out the excess water at the earliest</p> <p>Apply 4-5 kg N/acre after draining excess water</p> <p>Spray KNO<sub>3</sub> @1 % or water soluble fertilizers like 19-19-19 @ 1% to support nutrition</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>Drain out the excess water at the earliest</p> <p>Harvest the crop after the fields are dried up</p>

		and disease incidence		
Maize	<p>Drain out the excess water at the earliest</p> <p>Intercultivation and earthing up to be done</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>Drain out the excess water at the earliest</p> <p>Takeup weed control either mechanically or through weedicides</p> <p>Intercultivation and earthing up to be done</p> <p>Apply 20 kg N + 10 kg K/ha after draining excess water</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>Drain out the excess water at the earliest</p> <p>Take up plant protection measures against possible pests and disease incidence</p>	<p>Drain out the excess water at the earliest</p> <p>Cob picking to be done after they are dried fully</p>
<b>Horticulture crops – Fruits</b>				
Mango	<p>If the damage is severe, go for replanting.</p>	<p>Trees fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>Tress fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature fruits as soon as possible.</p> <p>Collect the fallen fruits and sell immediately or go for preparation of processed products.</p> <p>If to store, store the produce in well-ventilated place temporarily before it can be marketed.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>
Banana		<p>Wind damaged plants should be pruned using disinfected secaetures and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>The fallen tress may be cut leaving two suckers</p> <p>Inter-cultivate the soil with gorru</p>	<p>Wind damaged plants should be pruned using disinfected secaetures and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>The fallen tress may be cut leaving two suckers</p> <p>Topdressing of booster</p>	<p>Wind damaged plants should be pruned using disinfected secaetures and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible.</p> <p>Harvest the mature bunches as soon as possible. use ripening chambers for quick and uniform ripening</p> <p>Store the harvested bunches in well</p>



		<p>for aeration.  Spray 0.5 % KNO<sub>3</sub> or Urea 2% solution 2-3 times.  Topdressing of booster dose of 80 g MOP + 100 g Urea per plant at two to three times intervals.  Spray Propiconazole 1ml in one litre, 2-3 times against leaf spots.  Soil drenching with COC @ 3 g/litre to avoid rhizome rotting.  Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.  If the age of the plant is less than three months and submergence up to three feet better to replant the garden.</p>	<p>dose of 80 g MOP + 100 g Urea per plant at two to three times intervals  Mature bunches on the completely damaged plants be covered with Leaves and harvested with in 15-20days</p>	<p>ventilated place temporarily before it can be marketed.  Market the produce as soon as possible.  3-4 foliar application of KNO<sub>3</sub> on immature/developing bunches and leaves at weekly intervals.  Staking with bamboo for support</p>
Guava	<p>Drain the excess water as soon as possible  Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.  Provide support to the young plants.</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste  Drain the excess water as soon as possible  Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste  Drain the excess water as soon as possible  Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste  Drain the excess water as soon as possible.  Harvest the mature fruits as soon as possible.  Store the fruits in well-ventilated place temporarily before it can be marketed.  Market the fruits as soon as possible.  The unmarketable fruits may be utilized for processing</p>
Horticulture crops vegetables				
Chillies	<p>Grow nursery on raised beds.</p>	<p>Uprooted plants may be lifted and earthed up  Drain the excess water as soon</p>	<p>Uprooted plants may be lifted and earthed up  Drain the excess water as</p>	<p>Drain the excess water as soon as possible.  Dry the pods on concrete floor/</p>

		<p>as possible Gap filling must be done immediately If damage is more go for replanting Spray Urea 2% solution 2-3 times. Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>soon as possible Spray Urea 2% solution 2-3 times. Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>tarpaulins immediately Use poly house solar driers for quick drying Remove the pest and disease infected pods. .</p>
Tomato	<p>Grow nursery on raised beds. If damage is more go for resowing</p>	<p>Uprooted plants may be lifted and earthed up Drain the excess water as soon as possible Gap filling must be done immediately Spray Urea 2% solution 2-3 times. Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible. If damage is more ,go for replanting</p>	<p>Uprooted plants may be lifted and earthed up Drain the excess water as soon as possible Spray Urea 2% solution 2-3 times. Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible. Spray COC 30 g in 10 liters of water, 2-3 times against leaf spots. If damage is more ,go for replanting</p>	<p>Drain the excess water as soon as possible. Harvest the mature produce as soon as possible. Store the produce in well-ventilated place temporarily before it can be marketed. Market the produce as soon as possible.</p>
Cucumber		<p>Drain the excess water as soon as possible Spray Urea 2% solution 2-3 times. Topdressing of booster dose of 10 kg MOP + 30 kg Urea per acre as soon as possible. Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop. In case of severe damage (considered as complete</p>	<p>Drain the excess water as soon as possible Spray Urea 2% solution 2-3 times. Topdressing of booster dose of 10 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>Drain the excess water as soon as possible. Harvest the mature produce as soon as possible. Store the produce in well-ventilated place temporarily before it can be marketed. Market the produce as soon as possible.</p>

		economical loss), and the contingency period is between June to August, go for resowing		
Spices and Plantation crops				
Oil Palm Coconut	Planting should be done on mounts or bunds Drainage system suited to local conditions. may be provided to remove surplus water from root zone Relief drains [shallow] channels are opened at places where water accumulates and connected with main drain to remove water from the surface	Drain the excess water as soon as possible Twisted leaves may be cut and removed Apply booster dose of NPK fertilizers The palms have fallen with root system still having contact with the soil ,they need to be brought to position and provided with soil mound and support	Drain the excess water as soon as possible Hanging bunches may be provided with supports wherever possible .Apply booster dose of NPK fertilizers The palms have fallen with root system still having contact with soil they need to be brought to position and provided with soil mound and support	Twisted leaves may be cut and removed Hanging bunches may be provided with supports wherever possible Harvest the mature nuts as soon as possible. Market the produce as soon as possible.
Turmeric		Drain the excess water as soon as possible Spray Urea 2% or 1% KNO <sub>3</sub> followed by Ferrous Sulphate 0.5% + Citric Acid 0.1 % solution 2-3 times. Topdressing of booster dose of 40 kg MOP + 50 kg Urea along with 250 kg of Neem Cake per acre as soon as possible. In case of severe damage (considered as complete economical loss or if inundation is more than for four days), and the contingency period is between June to August, sowing of best alternative crop must be taken up.	Drain the excess water as soon as possible Spray Urea 2% or 1% KNO <sub>3</sub> followed by Ferrous Sulphate 0.5% + Citric Acid 0.1 % solution 2-3 times. Topdressing of booster dose of 40 kg MOP + 50 kg Urea along with 250 kg of Neem Cake per acre as soon as possible.	Drain the excess water as soon as possible. Harvest the rhizomes when field comes to normal Use boilers and polishers for processing Remove and separate the rotten and mould affected rhizomes. Cook and dry the rhizomes as soon as possible.

## 2.5 Contingent strategies for Livestock, Poultry & Fisheries

### 2.5.1.1 Livestock

Before the event	During the event	After the event
<b>Feed and fodder availability</b>		
<p>Conserving fodder/crop residues/ forest grass by silage / hay making either by individual or on community basis</p> <p>Preparing complete diets and storing in strategic locations</p> <p>Organize procurement of dry fodders / feed ingredients from surplus areas</p> <p>Establish fodder banks and feed banks</p> <p>Livestock relief camps during floods/cyclones must be planned in the vicinity of relief camps for people</p> <p>Capacity building and preparedness</p>	<p>Organise relief camps 2. Supply silage / hay to farmers with productive stock on subsidized rates</p> <p>Segregate old, weak and unproductive stock and send for slaughter</p> <p>Supply mineral mixture to avoid deficiencies</p> <p>Dry fodder must be offered to the livestock in little quantities for number of times</p> <p>Concentrate feed or complete feed must be offered to only productive and young stock only</p>	<p>Capacity building to stake holders on drought /cyclone/flood mitigation in livestock sector</p> <p>Promote fodder cultivation.</p> <p>Flushing the stock to recoup</p> <p>Avoid soaked and mould infected feeds / fodders to livestock</p> <p>Replenish the feed and fodder banks</p> <p>Promote fodder preservation techniques like silage / hay making</p>
<b>Drinking water</b>		
<p>Construct drinking water tanks in herding places, village junctions and in relief camp locations</p> <p>Plan for sufficient number of tanks for water transportation</p> <p>Identify bore wells, which can sustain demand.</p> <p>Procure sufficient quantities of water Sanitizers</p>	<p>Regular supply of clean drinking water to all tanks 2. Cleaning the tanks in regular intervals</p> <p>Keep the livestock away from contaminated flood/cyclone/stagnated waters</p> <p>Add water sanitizers</p>	<p>Hand over the maintenance of the structures to panchayats</p> <p>Sensitize the farming community about importance of clean drinking water</p>
<b>Health and disease Management</b>		

<p>Procure and stock emergency medicines and vaccines for important endemic diseases of the area</p> <p>All the stock must be immunized for endemic diseases of the area</p> <p>Carry out deworming to all young stock</p> <p>Keep stock of bleaching powder and lime</p> <p>Carry out Butax spray for control of external parasites</p> <p>Identify the Clinical staff and trained paravets and indent for their services as per schedules</p> <p>Identify the volunteers who can serve in need of emergency</p>	<p>Keep close watch on the health of the stock</p> <p>Sick animals must be isolated and treated Separately.</p> <p>Carry out deworming and spraying to all animals entering into relief camps</p> <p>Clean the animal houses regularly and apply disinfectants.</p> <p>Safe and hygienic disposal of dead animal carcasses</p> <p>Organize with community daily lifting of dung from relief camps</p>	<p>Keep close surveillance on disease outbreak.</p> <p>Undertake the vaccination depending on need</p> <p>Keep the animal houses clean and spray disinfectants</p>
---	---	--

	<b>Suggested contingency measures</b>		
	<b>Before the event</b>	<b>During the event</b>	<b>After the event</b>
Cyclone	<p>Harvest all the possible wetted grain (Rice/maize/blackgram/green gram etc) and use as animal feed.</p> <p>As the district is chronically prone for cyclone, arrange for storing minimum required quantity of hay (25-50 kg) and concentrates (10-25 kg) per animal in farmer's/LS keepers house/ shed for feeding during cyclone.</p> <p>Stock of anti-diarrheal drugs and electrolytes should be made available for emergency transport</p> <p>Don't allow the animals for grazing in case of early forewarning (EFW) of cyclone</p> <p>Incase of EFW of severe cyclone, shift the animals to safer places.</p>	<p>Treatment of the sick, injured and affected animals through arrangement of mobile emergency veterinary hospitals / rescue animal health workers.</p> <p>Diarrhea out break may happen. Health camps should be organized</p> <p>In severe cases un-tether <b>or</b> let loose the animals</p> <p>Arrange transportation of highly productive animals to safer place</p> <p>Spraying of fly repellants in animal sheds</p>	<p>Repair of animal shed</p> <p>Deworm the animals through mass camps</p> <p>Vaccinate against possible disease out breaks like HS, BQ, FMD and PPR</p> <p>Proper dispose of the dead animals / carcasses by burning / deep burying (4-8 feet) with lime powder (1kg for small ruminants and 5kg for large ruminants) in pit</p> <p>Bleach / chlorinate (0.1%) drinking water or water resources</p> <p>Collect drowned crop material, dry it and store for future use</p> <p>Sowing of short duration fodder crops in unsown and water logged areas when crops are damaged and no chance to replant</p> <p>Application of urea (20-25kg/ha) in</p>

			the inundated areas and CPR's to enhance the bio mass production.
Floods	<p>In case of early forewarning (EFW), harvest all the crops (Rice/maize/blackgram/green gram) that can be useful as fodder in future (store properly)</p> <p>Don't allow the animals for grazing if severe floods are forewarned</p> <p>As regularly flood prone district, arrange for storing minimum required quantity of hay (25-50kg) and concentrates (25kgs) per animals in farmer / LS keepers house / shed for feeding animals during floods</p> <p>Arrangement for transportation of animals from low lying area to safer places and also for rescue animal health workers to get involve in rescue operations</p>	<p>Transportation of animals to elevated areas</p> <p>Stall feeding of animals with stored hay and concentrates</p> <p>Proper hygiene and sanitation of the animal shed</p> <p>In severe floods, un-tether or let loose the animals</p> <p>Emergency outlet establishment for required medicines or feed in each village</p> <p>Spraying of fly repellants in animal sheds</p>	<p>Repair of animal shed</p> <p>Bring back the animals to the shed</p> <p>Cleaning and disinfection of the shed</p> <p>Bleach (0.1%) drinking water / water sources</p> <p>Deworming with broad spectrum dewormers</p> <p>Vaccination against possible disease out breaks like HS, BQ, FMD and PPR</p> <p>Proper disposable of the dead animals / carcasses by burning / deep burying (4-8 feet) with lime powder (1kg for small ruminants and 5kg for large ruminants) in pit</p> <p>Drying the harvested crop material and proper storage for use as fodder.</p>
Health and Disease management	<p>List out the endemic diseases (species wise) in that district and store vaccines for those diseases</p> <p>Timely vaccination (as per enclosed vaccination schedule) against all endemic diseases</p> <p>Surveillance and disease monitoring network to be established at Joint Director (Animal Husbandry) office in the district</p>	<p>Constitution of Rapid Action Veterinary Force</p> <p>Procurement of emergency medicines and medical kits</p> <p>Rescue of sick and injured animals and their treatment</p>	<p>Conducting mass animal health camps</p> <p>Conducting fertility camps</p> <p>Mass deworming camps</p>
Insurance	Encouraging insurance of livestock	Listing out the details of the dead animals	<p>Submission for insurance claim and availing insurance benefit</p> <p>Purchase of new productive animals</p>
Drinking water	<p>Identification of water resources</p> <p>Rain water harvesting and create water bodies/watering points (when water is scarce use only as drinking water for animals)</p>	Restrict wallowing of animals in water bodies/resources	<p>Bleach (0.1%) drinking water / water sources</p> <p>Provide clean drinking water</p>

Drought			
Feed & Fodder availability	<p>Available paddy straw and sugar cane tops should be properly stored for future use.</p> <p>Chopping of fodder should be made as mandatory in every village through supply and establishment of good quality chaff cutters.</p> <p>Harvesting and collection of perennial vegetation particularly grasses which grow during monsoon</p> <p>Proper drying, baling and densification of harvested grass from previous season</p> <p>Creation of permanent fodder, feed and fodder seed banks in all drought prone areas</p>	<p>Harvest and use biomass of dried up crops (Rice/maize/greengram/blackgram) material as fodder.</p> <p>Harvest the tree fodder (Neem, Subabul, Acasia, Pipal etc) and unconventional feeds resources available and use as fodder for livestock (LS).</p> <p>Available feed and fodder should be cut from CPRs and stall fed in order to reduce the energy requirements of the animals</p> <p>Hay should be transported to the needy areas from the near by districts in case of drought</p> <p>Advise the farmers about the practice of mixing available kitchen waste with dry fodder while feeding</p>	<p>Short duration fodder crops of sorghum/bajra/maize(UP chari, MP chari, HC-136, HD-2, GAIN T BAJRA, L-74, K-677, Ananad/African Tall, Kisan composite, Moti, Manjari, B1-7 should be sown in unsown and crop failed areas where no further routine crop sowing is not possible</p>

#### Vaccination programme for cattle and buffalo

Disease	Age and season at vaccination
Anthrax	In endemic areas only, Feb to May
Haemorrhagic septicaemia (HS)	May to June
Black quarter (BQ)	May to June
Foot and mouth disease (FMD)	July/August and November/December

#### Vaccination schedule in small ruminants (Sheep & Goat)

Disease	Season
Foot and mouth disease (FMD)	Preferably in winter / autumn
Peste des Petits Ruminants (PPR)	Preferably in January
Black quarter (BQ)	May / June
Enterotoxaemia (ET)	May
Haemorrhagic septicaemia (HS)	March / June

Sheep pox (SP)	November
----------------	----------



## 2.5.2 Poultry

	Suggested contingency measures		
	Before the event	During the event	After the event
<b>Drought</b>			
Shortage of feed ingredients	Storing of house hold grain like maize, broken rice etc, in to use as feed in case of severe drought.	Supplementation only for productive birds with house holds grain. Supplementation of shell grit (calcium) for laying birds. Culling of weak birds.	Supplementation to all survived birds.
Drinking water		Use water sanitizers or offer cool drinking water	
Health and disease management	Culling of sick birds. Deworming and vaccination against RD and fowl pox	Mixing of Vit. A,D,E, K and B-complex including vit C in drinking water (5ml in one litre water).	Hygienic and sanitation of poultry house. Disposal of dead birds by burning / burying with lime powder in pit.
<b>Floods</b>			
Shortage of feed ingredients	In case of early forewarning of floods, shift the birds to safer place. Storing of house hold grain like maize, broken rice, bajra etc,	Use stored feed as supplement. Don't allow for scavenging Culling of weak birds.	Routine practices are followed Deworming and vaccination against RD .
Drinking water		Use water sanitizers or offer cool drinking water.	
Health and disease management	In case of EFW, add antibiotic powder (Terramycin/Ampicilline/ Ampiclox etc., 10g in one litre) in drinking water to prevent any disease outbreak	Prevent water logging surrounding the sheds through proper drainage facility. Assure supply of electricity by generator or solar energy or biogas. Sprinkle lime powder to prevent ammonia accumulation due to dampness.	Sanitation of poultry house. Treatment of affected birds. Disposal of dead birds by burning / burying with lime powder in pit . Disposal of poultry manure to prevent protozoal problem Supplementation of

			coccidiostats in feed. Vaccination against RD.
<b>Cyclone</b>			
Shortage of feed ingredients	In case of EFW, shift the birds to safer place. Storing of house hold grain like maize, broken rice, bajra etc., Culling of weak birds.	Use stored feed as supplement Don't allow for scavenging Protect from thunder storms	Routine practices are followed
Drinking water		Use water sanitizers or offer cool drinking water	
Health and disease management	In case of EFW, add antibiotic powder in drinking water to prevent any disease outbreak.	Sanitation of poultry house. Treatment of affected birds. Prevent water logging surrounding the sheds Assure supply of electricity. Sprinkle lime powder (5-10g per square feet) to prevent ammonia accumulation due to dampness.	Disposal of dead birds by burning / deep burying with lime powder in pit. Disposal of poultry manure to prevent protozoal problem. Supplementation of coccidiostats in feed Vaccination against Ranikhet Disease (0.5ml S/c).

### 2.5.3 Fisheries/ Aquaculture

	Suggested contingency measures		
	Before the event	During the event	After the event
<b>1) Drought</b>			
<b>A. Capture</b>			
Marine	No intervention	No intervention	No intervention
Inland			
(i) Shallow water depth due to insufficient rains/inflow	Stocking of advanced fingerlings in half or even less than the normal stocking density or stocking of common carp seed	Immediate harvesting or decreasing the density commensurate with the water quantity.	De weeding and deepening of tank to ensure retention of water for a longer period and provision of employment under MGNREGP
(ii) Changes in water quality	Regular monitoring of water quality parameters and application of geolites, soil probiotics, etc to maintain water quality	Immediate harvesting or changing the water quality by application of sanitisers.	Removal of top layer, deep ploughing of tank and application of lime
(iii) Any other			
<b>B. Aquaculture</b>			
(i) Shallow water in ponds due to insufficient rains/inflow	Crop holiday or going for stocking of yearlings by reducing the density according to availability of water	Harvesting of fish and leaving the pond fallow till next season	Removal of top layer, deep ploughing of tank and application of lime
(ii) Impact of salt load build up in ponds / change in water quality	Stocking of salinity tolerant fish / shrimp, application of geolites and other buffers	Frequent change of water with fresh water	Frequent draining of the pond with fresh water, removal of top layers
(iii) Any other			
<b>2) Floods</b>			
<b>A. Capture</b>			
Marine	No intervention	No intervention	No intervention
Inland			
(i) Average compensation paid due to loss of human life	Shifting the people from low lying areas to relief camps	Deployment of specially trained persons for rescue operations by providing life bouys, jackets, ropes, boats, etc	Payment sufficient ex-gratia to the families

(ii) No. of boats / nets/damaged	Shifting and relocating boats and nets to safer places when warnings are issued, to avoid fishing, etc	Shifting and relocating boats and nets to safer places	Assessment of damages to boats and nets and provision of boats and nets for restoration of livelihoods
(iii) No. of houses damaged	Avoidance of construction of houses in flood prone areas, construction of pucca houses at elevated places,	Shifting of people by relief boats to the relief camps	Assessment of damages to houses and provision of compensation in case of partial damage and sanction house under existing schemes
(iv) Loss of stock	Avoidance of surface species like catla, silver carp since they are vulnerable in tanks prone to floods, erection of nets across the spill way or just beyond it	Erection of nets at spill ways	Taking up compensatory stocking
(v) Changes in water quality		When dissolved oxygen levels go down, aerators, recirculation of water, etc are to be attempted to maintain DO levels, going for partial harvest, etc	
(vi) Health and diseases	Sometimes there may be heavy accumulation of nutrients and organic matter.	There may be break out of Hemorrhagic septicemia. Addition of antibiotics like Chloro Tetra Cycline or Oxy Tetra Cycline to the feed to control the disease	Removal of weeds, top layer of soil, deep ploughing of tank and application of lime, exposing to sun light
<b>B. Aquaculture</b>			
(i) Inundation with flood water	Raising and rivetting the bunds, construction of spill way to release excess water, erection of nets to avoid escape of fish	Continuous pumping of excess water, erection of nets low lying areas	Strengthening of bunds, excavating channels along the sides of the ponds for free escape of water
(ii) Water continuation and changes in water quality		When dissolved oxygen levels go down, aerators, recirculation of water, etc are to be attempted to maintain DO levels, going for partial harvest, etc	
(iii) Health and diseases	Sometimes there may be heavy accumulation of nutrients and organic matter.	There may be break out of Hemorrhagic septicemia. Addition of antibiotics like Chloro Tetra Cycline or Oxy Tetra Cycline to the feed to control the disease	Removal of weeds, top layer of soil, deep ploughing of tank and application of lime, exposing to sun light

(iv) Loss of stock and inputs (feed, chemicals etc)	Advance erection of nets, strengthening of bunds where they are prone to breaches, harvesting or reducing the density	Suspension of feeding, application of organic manures	Compensatory stocking, assessment of values and payment of subsidy on inputs
(v) Infrastructure damage (pumps, aerators, huts etc)	Insuring pond, accessories, etc., Shifting of aerators, pumps soon after warnings are issued	Relocating pumps, aerators to elevated places	Assessment of damages and provision of them on subsidy
(vi) Any other			
<b>3. Cyclone / Tsunami</b>			
A. Capture			
Marine			
(i) Average compensation paid due to loss of fishermen lives	Avoidance of fishing, preventing fishermen from venturing into sea, carrying of safety equipment and VHF sets, shifting fishermen from vulnerable areas to relief camps, etc	To ensure the return of fishing boats on long voyages, provision of information on such boats to coast Guard	Payment sufficient ex-gratia to the families
(ii) Avg. no. of boats / nets/damaged	Avoidance of fishing when warnings are issued, shifting of boats and nets to safe places	Shifting and relocating boats and nets to safer places	Assessment of damages to boats and nets and provision of boats and nets for restoration of livelihoods
(iii) Avg. no. of houses damaged	Avoidance of houses in Coastal Regulation Zone, designing of houses to withstand impact of turbulent wind and water	Shifting of people by relief boats to the relief camps	Assessment of damages to houses and provision of compensation in case of partial damage and sanction house under existing schemes
Inland	Erection of protective nets across the surplus weir to prevent fish loss due to overflows	Continuous monitoring to prevent or minimise escape of fish along with surplus water	Compensatory stocking of seed
B. Aquaculture			
(i) Overflow / flooding of ponds	The design of the pond must be in such a manner as to bail out surplus water and to prevent loss of standing crop	Continuous monitoring to prevent or minimise escape of fish along with surplus water	Compensatory stocking of seed
(ii) Changes in water quality (fresh water / brackish water ratio)	Recirculation water to replenish and ensure sufficient dissolved oxygen levels in the pond. Maintenance of	Continuation of the same process.	Restoration of physical and chemical parameters

	salinity levels by pumping in water from creeks.		
(iii) Health and diseases	Removal of stress causing factors to maintain the health of the animal	Removal of stress causing factors to maintain the health of the animal	Restoration of physical and chemical parameters
(iv) Loss of stock and inputs (feed, chemicals etc)	Preventive nets must be erected to minimise loss of stock	Continuation of the same process.	Compensatory stocking of seed
(v) Infrastructure damage (pumps, aerators, shelters/huts etc)	Pumps, aerators, etc must be protected by moving them to safe locations	To avoid use of aerators, pumps and other appliances	Overhauling of the equipment to prevent from being damaged
(vi) Any other			
<b>4. Heat wave and cold wave</b>			
<b>A. Capture</b>			
Marine	Avoidance of fishing	Avoidance of fishing	No intervention
Inland	Monitoring dissolved oxygen levels	Monitoring dissolved oxygen levels	No intervention
<b>B. Aquaculture</b>			
(i) Changes in pond environment (water quality)	Reduction of biomass by partial harvest in the event of heat as the DO levels will be very low.	Avoidance of fishing	Compensatory stocking of seed and restoration of all physical and chemical parameters
(ii) Health and Disease management	Removal of stress causing factors to maintain the health of the animal	Removal of stress causing factors to maintain the health of the animal	Compensatory stocking of seed and restoration of all physical and chemical parameters
(iii) Any other			

# Contingency Plans for Rabi Crops for Krishna Zone

## Krishna District

### 1. Rainfall Information

	Oct – Dec	Jan – Mar
(a) Normal rainfall during <i>Rabi</i> season :	250	-
(b) Number of rainy days :	7-10	-

### 2. Rabi crops cultivated

#### 2a Area Production statistics

S. NO	Name of the crop	Area ( ha)	Productivity ( Kg/ha)	Production (Lakh MTs)
		Kharif 2018	Kharif 2018	Kharif 2018
1	Rice	245815	6118	15.04
2	Maize	4063	4516	0.18
3	Redgram	1972	874	1.017
4	Greengram	4158	970	0.04
5	Blackgram	1972	1039	0.02
6	Groundnut	1029	3207	0.03
7	Cotton	47109	1878	0.88

8	Sugarcane	13132	99587	13.08
---	-----------	-------	-------	-------

## 2b Source wise (Water) cultivated area

Sources of Irrigation	Number	Area ('000 ha)	Percentage of total irrigated area
Canals	KE canal System	251	73.82
Tanks	983	38	11.18
Open wells	15552		
Bore wells	13549	30	8.82
Lift irrigation			
Micro-irrigation			
Other sources	64	21	6.18
Total Irrigated Area		340	100.0

## 3. Sowing window information

S. No.	Soil type	Cropping system	Crop name	Optimum sowing window
1	Deltaic black soils	Rice-pulse	Blackgram/greengram	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of December
2		Rice-Maize	Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January
3		Rice-Rice	Rice	2 <sup>nd</sup> fortnight of November to 1 <sup>st</sup> Fortnight of December
4	Sandy loams	Rice-Groundnut	Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December
5	Black soils	Pulse-Bengalgram	Bengalgram	October 15 <sup>th</sup> to November 15 <sup>th</sup>



6	Black/Red soils	Pulse-Maize/ Maize-Maize	Maize	1 <sup>st</sup> FN of October to 1 <sup>st</sup> FN of November
7		Pulse-Pulse	Blackgram	3 <sup>rd</sup> week of October to 2 <sup>nd</sup> week of November

#### 4. Contingency measures Field crops

**For crops grown with residual moisture i.e., under rainfed condition**

(a) Excess residual moisture

S. No.	Soil type	Cropping system	Crop name	Sowing Window	Variety	Management practices
1	Deltaic black soils/Red and black soils	Rice-pulse	Blackgram	3 <sup>rd</sup> week of November to 3 <sup>rd</sup> week of December	LBG-752 LBG-787 PU-31, GBG-1, TBG-104	Foliar spray of urea 2% or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> and FeSO <sub>4</sub> spray; Spraying of copper oxy chloride or mancozeb etc to manage fungal diseases.
2			Greengram		LGG 460 TM 96-2 WGG 42 IPM 2-14	

3			Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January	DHM-115, 117, KH-510 Private hybrids	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> and FeSO <sub>4</sub> spray; Spraying of copper oxy chloride or mancozeb etc. to manage fungal diseases. Intercultivation to reduce soil moisture
4	Sandy loams		Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K6, K9, Kadiri Harithandra, Dharani	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> and FeSO <sub>4</sub> spray; Spraying of copper oxy chloride or mancozeb etc. to manage fungal diseases.

(b) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

S. No	Soil type	Cropping system	Crop name	Sowing Window	Variety	Management practices
1	Sandy loams	Rice-Groundnut	Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, gypsum application, mulching to conserve moisture, need based application of

						pesticides to manage insect pests and diseases
2	Black soils	Pulse-Bengalgram	Bengalgram	October 15 <sup>th</sup> to November 15 <sup>th</sup>	JG-11, NBeG-47, 49	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, mulching to conserve moisture, need based application of pesticides to manage insect pests and diseases
3	Black/ Red soils	Pulse-Maize/ Maize-Maize	Maize	1 <sup>st</sup> FN of October to 1 <sup>st</sup> FN of November	DHM 113, Private hybrids	Foliar spray of urea 2% or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, mulching to conserve moisture, control of fall army worm, stem borers, tursicum blight, sheath blight etc.,
4		Pulse-Pulse	Blackgram	3 <sup>rd</sup> week of October to 2 <sup>nd</sup> week of November	PU-31, TBG 104	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, need based application of

						pesticides to manage insect pests and diseases i.e. Maruca pod borer, sucking insect pests, Spodoptera, Powdery mildew etc.
--	--	--	--	--	--	---

(c) Severe limitation in moisture. Deficit of rainfall during September/October months by more than 40%.

S. No	Soil type	Cropping system	Crop name	Sowing Window	Variety	Management practices
1	Sandy loams	Rice-Groundnut	Blackgram	3 <sup>rd</sup> week of October to 2 <sup>nd</sup> week of November	PU-31, TBG 104	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, need based application of pesticides to manage insect pests and diseases i.e. Maruca pod borer, sucking insect pests, Spodoptera, Powdery mildew etc.
2	Black soils	Pulse-Bengal gram	Bengal gram	October 15 <sup>th</sup> to November 15 <sup>th</sup>	JG-11, NBeG-47, 49	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, mulching to

						conserve moisture, need based application of pesticides to manage insect pests and diseases
3	Black/ Red soils	Pulse-Maize/ Maize-Maize	Blackgram	3 <sup>rd</sup> week of October to 2 <sup>nd</sup> week of November	PU-31, TBG 104	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, need based application of pesticides to manage insect pests and diseases i.e. Maruca pod borer, sucking insect pests, Spodoptera, Powdery mildew etc.
4		Pulse-Pulse	Blackgram	3 <sup>rd</sup> week of October to 2 <sup>nd</sup> week of November	PU-31, TBG 104	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> spray 2-3 times at regular intervals, need based application of pesticides to manage insect pests and diseases i.e. Maruca pod borer, sucking insect pests, Spodoptera, Powdery mildew etc.

**For crops grown with groundwater**

(a) Above normal rainfall in *Kharif* coupled with good distribution

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Red/black	Pulse/Maize-Maize	Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January	DHM-115, 117, KH-510 Private hybrids	Pre and post emergence application of weedicides, Control of fall army worm, stem borers, tursicum blight, sheath blight etc., correction of nutrient deficiencies, irrigation at critical growth stages or based on soil moisture condition
2	Sandy loams	Rice-Groundnut	Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
3	Black soils	Pulse/Fallow-Bengalgram	Bengalgram	October 15 <sup>th</sup> to November 15 <sup>th</sup>	JG-11, NBeG-47, 49	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to

						manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
--	--	--	--	--	--	--

(b) Normal rainfall

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Red/black	Pulse/Maize-Maize	Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Normal practices and need based plant protection measures
2	Sandy loams	Rice-Groundnut	Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
3	Black soils	Pulse/Fallow-Bengalgram	Bengalgram	October 15 <sup>th</sup> to November 15 <sup>th</sup>	JG-11, NBeG-47, 49	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and

						diseases, irrigation at critical growth stages or based on soil moisture condition
--	--	--	--	--	--	--

(c) Deficient rainfall in *Kharif* season (25-50% deficient)

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Red/ black	Pulse/Maize-Maize	Fodder Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Normal practices and need based plant protection measures
			Horsegram	4 <sup>th</sup> week of Sept	CRIDA 18-R CRIDA-22B	Normal practices and need based plant protection measures
2	Sandy loams	Rice-Groundnut	Cowpea	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TPTC-29, C 152, CO-8, APFC-10/1	Normal practices and need based plant protection measures
3	Black soils	Pulse/Fallow-Bengalgram	Bengalgram	October 15 <sup>th</sup> to November 15 <sup>th</sup>	JG-11, NBeG-47, 49	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition



(d) Scanty rainfall in *Kharif* season

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Red/ black	Pulse/Maize- Maize	Fodder Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Normal practices and need based plant protection measures
			Horsegram	4 <sup>th</sup> week of Sept	CRIDA 18-R CRIDA-22B	Normal practices and need based plant protection measures
2	Sandy loams	Rice-Groundnut	Cowpea	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TPTC-29, C 152, CO-8, APFC-10/1	Normal practices and need based plant protection measures
3	Black soils	Pulse/fallow- Bengalgram	Bengalgram	October 15 <sup>th</sup> to November 15 <sup>th</sup>	JG-11, NBeG-47, 49	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition

## Management practices for unseasonal rains

Condition - Continuous high rainfall in a short span leading to water logging and Heavy rainfall with high speed winds in a short span				
Crop	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Rice	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>• Take up gap filling either with available nursery or by splitting the tillers from the surviving hills</li> <li>• Take up suitable plant protection Measures in anticipation of pest &amp; disease out breaks</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>• Take up suitable plant protection Measures in anticipation of pest &amp; disease out breaks</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Take up suitable plant protection measures in anticipation of pest &amp; disease out breaks</li> </ul>	<ul style="list-style-type: none"> <li>• Drain out water and spread sheaves loosely in field or field bunds where there is no water stagnation</li> <li>• Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds</li> </ul> <p>Thresh after drying the sheaves properly Ensure proper grain moisture before storing</p>
Blackgram/ Greengram/ Bengal gram	<p>Drain the excess water as early as possible</p> <p>Apply 4-5 kg N /ha after draining excess water</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, at 1% to support nutrition</p> <p>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb</p>	<p>Drain the excess water as early as possible</p> <p>Apply 4-5 kg N /ha after draining excess water</p> <p>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition</p> <p>Spray fungicides like Copper oxy</p>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Allow the crop to dry completely before harvesting</li> </ul>	<ul style="list-style-type: none"> <li>• Spread the bundles drenched in rain on field bunds or drying floors to quicken the drying</li> <li>• Thresh the bundles after they are dried properly Dry the grain to proper <ul style="list-style-type: none"> <li>• moisture per cent before</li> <li>• bagging and storing to prevent</li> </ul> </li> </ul>

	<p>0.25% two to three times by rotating the chemicals</p> <ul style="list-style-type: none"> <li>Take up timely control measures against the outbreak of pests like Spodoptera etc.</li> </ul>	<p>chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</p> <p>Take up timely control measures against the outbreak of pests like Spodoptera etc.</p>		deterioration in quality during storage
Maize	<ul style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>Take up inter cultivation and at optimum soil moisture condition to loosen and aerate the soil and to control weeds</li> <li>Earthen up the crop for anchorage Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition</li> <li>Take up timely control measures for Pink stem borer, sheath blight and Turicum leaf blight</li> </ul>	<ul style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>Spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19 at 1% to support nutrition Take up timely control measures for sheath blight and post flowering</li> <li>stalk rots</li> </ul>	<p>Drain the excess water as early as possible</p> <p>Allow the crop to dry completely before harvesting</p>	<ul style="list-style-type: none"> <li>Harvest the cobs after the they are dried up properly. Dry the grain to optimum moisture condition before storing</li> </ul>

**For crops grown with Canal Irrigation: The scenario would be based on the storage available in the reservoirs.**

a. Limited release of water

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soil & Red soils	Rice – Rice	Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January	DHM-115, 117, KH-510 Private hybrids	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> and FeSO <sub>4</sub> spray; Spraying of copper oxy chloride or mancozeb etc. to manage fungal diseases. Intercultivation to reduce soil moisture
			Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup>	NLR-34449	Direct sowing with drum seeder

			Week of Nov	NLR-3042 NLR-4001 JGL-384 MTU-1156 MTU-1153 MTU-1121	
2			Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,
3			Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-399
4			Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani
5			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752
					Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
					Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition

b. Delayed release of water

**Head reach:**

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soils/Red soils	Rice-Rice	Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	NLR-34449 NLR-3042 NLR-4001 JGL-384 MTU-1156 MTU-1153 MTU-1121	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition

**Middle reach:**

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soils/Red soils	Rice-Rice	Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	NLR-34449 NLR-3042 NLR-4001 JGL-384 MTU-1156	Direct sowing with drum seeder

					MTU-1153	
					MTU-1121	

**Tail end:**

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soil & Red soils	Rice – Rice	Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
2			Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-399	Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
3			Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	
			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on

						soil moisture condition
--	--	--	--	--	--	-------------------------

**5. Contingency measures for Horticulture Crops (Existing / New plantations)**

S. No.	Crop Name	Specific management practices to be taken up following excess/deficient/scanty rainfall	Time of intervention	Remarks

**6. Contingency measures for Horticulture Crops (vegetables)**

S. No.	Crop Name	Specific management practices to be taken up following excess/deficient/scanty rainfall	Time of intervention	Remarks

**7. Temperature related stresses for field and horticulture crops:**

Excess temperatures/ Less than normal temperatures

S.No	Crop name	Stage of crop growth	Threshold temperature*	Suggested management practices


\* Temperature increase or decrease over normal and for number of days. For example, increase of 3 degrees over normal for a period of 5 days

**8. Management practices for livestock** (to cover shelter management during cold or heat waves, production/regulation of fodder in rabi season in deficient monsoon years/ excess monsoon rainfall years etc),

**For Fodder crops grown with residual moisture i.e., under rainfed condition**

(a) Excess (rainfall during September/October months) residual moisture

S. No.	Soil type	Cropping system	Fodder name	Variety	Management practices

(b) Normal rainfall (rainfall during September/October months) residual moisture

S. No.	Soil type	Cropping system	Crop name	Variety	Management practices



(b) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

S. No.	Soil type	Cropping system	Fodder name	Variety	Management practices

(c) Severe limitation in moisture. Deficit of rainfall during September/October months by more than 40%.

S. No.	Soil type	Cropping system	Fodder name	Variety	Management practices

**For fodder crops (mostly perennial fodder varieties as sole fodder crop) grown with groundwater**

S. No.	Soil type	Fodder name	Variety	Management practices

**Livestock management during severe cold waves/heat waves**

Nutritional management	Shelter management	Health management	Miscellaneous, if any