

**State: ANDHRA PRADESH**

**Agriculture Contingency Plan for District: PRAKASAM**

1.0 District Agriculture profile					
1.1	Agro-Climatic/Ecological Zone				
	Agro Ecological Sub Region (ICAR)	Eastern Coastal plain, hot sub-humid to semi arid eco region (7.3, 18.3)			
	Agro-Climatic Region (Planning Commission)	East Coast plain and hill region (XI)			
	Agro Climatic Zone (NARP)	Krishna Zone (AP-1)			
	List all the districts or part thereof falling under the NARP Zone	Guntur, Krishna, Prakasam			
	Geographic coordinates of district	Latitude	Longitude		Altitude
		14°57’N	78°43’E		224 m
	Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/RRTTS	Regional agricultural Research Station, Lam, Guntur			
	Mention the KVK located in the district	Darsi, Prakasam Dt , AP			
1.2	Rainfall	Normal RF (mm)	Normal Rainy days (no)	Normal Onset ( specify week and month)	Normal Cessation (specify week and month)
	SW monsoon (June-Sep):	388	36	June 2 <sup>nd</sup> week	October 2 <sup>nd</sup> week
	NE Monsoon(Oct-Dec):	393	18	October 2 <sup>nd</sup> week	Last week of December
	Winter (Jan- February)	16	5	-	-
	Summer (Mar-May)	73	6	-	-
	Annual	870	65	-	-

<b>1.3</b>	<b>Land use pattern of the district</b> (latest statistics)	<b>Geographical Area</b>	<b>Forest area</b>	<b>Land under non-agricultural use</b>	<b>Permanent pastures</b>	<b>Cultivable wasteland</b>	<b>Land under Misc. tree crops and groves</b>	<b>Barren and uncultivable land</b>	<b>Current fallows</b>	<b>Other fallows</b>
	<b>Area ('000 ha)</b>	1762.6	442.5	171.5	58.6	69.6	11.1	158.3	100.4	103.9

<b>1.4</b>	<b>Major Soils (common names like shallow red soils etc.,)</b>	<b>Area ('000 ha)</b>	<b>Percent (%) of total</b>
	1. Shallow Red soils	215	51
	2. Deep black cotton soils	173	41
	3. Sandy loamy soils	25	6
	4. Sandy soils	9	2
	Others (specify):		
<b>1.5</b>	<b>Agricultural land use</b>	<b>Area ('000 ha)</b>	<b>Cropping intensity %</b>
	Net sown area	592.5	107.7
	Area sown more than once	20.3	
	Gross cropped area	605.1	

1.6	Irrigation	Area ('000 ha)		
	Net irrigated area	173.1		
	Gross irrigated area	188.3		
	Rainfed area	375.0		
	Sources of Irrigation	Number	Area ('000 ha)	Percentage of total irrigated area
	Canals		66.7	34.3
	Tanks	957	27.8	14.3
	Open wells	22783		
	Bore wells	41163	84.4	43.4
	Lift irrigation	1407		
	Micro-irrigation	--		
	Other sources	--	15.7	8.1
	Total Irrigated Area		194.6	100.0
		--		
	Pump sets	--	--	
	No. of Tractors	--	--	
	Groundwater availability and use* (Data source: State/Central Ground water Department /Board)	No. of blocks/ Tehsils	(%) area	
	Over exploited			
	Critical			
	Semi- critical			
	Safe			
	Wastewater availability and use			
	Ground water quality			

\*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%

1.7		Major Field Crops cultivated	Area ('000 ha)					
			<i>Kharif</i>		<i>Rabi</i>		Summer	Total
			<i>Irrigated</i>	<i>Rainfed</i>	<i>Irrigated</i>	<i>Rainfed</i>		
	1	Chick Pea	-	-	153.0	-	-	153.0
	2	Paddy	32.00	--	39.00	--	--	71.0
	3	Redgram		63.60		0.30	--	63.9
	4	Tobacco				63.4		63.4
	5	Cotton	--	58.2	--	--	10.20	68.4
	6	Sesamum	--	8.12				
	7	Bajra		17.70	0.58	-	--	18.3
	8	Groundnut	1.5	--	2.80	--	--	4.3
	9	Castor	--	4.50	1.60	--	--	6.10
	10	Bengalgram	--	--	--	81.89	--	81.89
	11	Blackgram	--	4.72	--	20.90	--	25.62
	12	Chillies	23.89	--	10.88	--	--	34.77
	13	Other crops	22.8	--	2.5	--	--	25.3
		<b>Horticulture crops - Fruits</b>	<b>Total area</b>					
	1	Orange & Batavia	22.5					
	2	Mango	6.6					
	3	Sapota	5.6					
	4	Lemon	2.3					
	5	Papaya	1.6					
		<b>Horticultural crops - Vegetables</b>	<b>Total area</b>					
	1	Chillies	10.6					
	2	Tomato	6.9					
		<b>Plantation crops</b>	<b>Total area</b>					
	1	Coriander	1.3					

1.8	Livestock	Male ('000)	Female ('000)	Total ('000)
	Non descriptive Cattle (local low yielding)	77.3	35.1	112.4
	Crossbred cattle	1.4	1.0	2.4
	Non descriptive Buffaloes (local low yielding)	195.9	1077.9	1273.8

	Graded Buffaloes						
	Goat			436.5			
	Sheep			1478.6			
	Others (Camel, Pig, Yak etc.)			19.91			
	Commercial dairy farms (Number)						
1.9	Poultry	No. of farms	Total No. of birds ('000)				
	Commercial		293.0				
	Backyard		110.8				
1.10	Fisheries (Data source: Chief Planning Officer)						
	A. Capture						
	i) Marine (Data Source: Fisheries Department)	No. of fishermen	Boats		Nets	Storage facilities (Ice plants etc.)	
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Gill nets)		Non-mechanized (Shore Seines, Stake & trap nets)
		12748	38	937 / 2260	31 / 83789	0 / 807	20 / 1
	ii) Inland (Data Source: Fisheries Department)	No. Farmer owned ponds		No. of Reservoirs		No. of village tanks	
		70		9		159	
	B. Culture						
		Water Spread Area (ha)		Yield (t/ha)	Production ('000 tons)		
	i) Brackish water (Data Source: MPEDA/ Fisheries Department)	2380		0.001	2.731		
	ii) Fresh water (Data Source: Fisheries Department)	341		0.003	0.943		
	Others			0.000	27.215		

1.11	Production and Productivity of major crops (Average of last 5 years)	Kharif		Rabi		Summer		Total		Crop residue as fodder ('000 tons)
		Production (MT )	Productivity (kg/ha)	Production (MT )	Productivity (kg/ha)	Production (MT )	Productivity (kg/ha)	Production (MT )	Productivity (kg/ha)	
Major Field crops (Crops to be identified based on total acreage)										

[illegible]

1.12	Sowing window for 5 major field crops (start and end of normal sowing period)	Paddy	Redgram	Cotton	Bengal gram	Blackgram
	Kharif- Rainfed	--	June last week to August 2 <sup>nd</sup> week	June last week to August 2 <sup>nd</sup> week	--	June 2nd week to July last week
	Kharif-Irrigated	August 2 <sup>nd</sup> week to September 3 <sup>rd</sup> week	--	--	--	--
	Rabi- Rainfed	--	October 2nd week	--	October last week to November last week	September 2nd week to October 2 <sup>nd</sup> week
	Rabi-Irrigated	October 1 <sup>st</sup> week to December 1 <sup>st</sup> week	--	--		


1.13	What is the major contingency the district is prone to? (Tick mark and mention years if known during the last 10 year period)	Regular	Occasional	None
1	Drought	√		
2	Flood		√	
3	Cyclone		√	
4	Hail storm			
5	Heat wave			
6	Cold wave			
7	Frost			
8	Sea water intrusion			
9	Pests and diseases (specify)	<u>Rice</u> : Blast, BLB, BPH <u>Redgram</u> : SMD, Maruca and wilt <u>Cotton</u> : Pink boll worm, Sucking pest complex		

		<u>Castor</u> : Botrytis grey mould, Semilooper <u>Blackgram</u> : YMV <u>Bengalgram</u> : Wilt, Spodoptera exigua <u>Maize</u> : Fall army worm, leaf blight		
10	Others (Fog)		√	

1.14	Include Digital maps of the district for	Location map of district within State as Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure 2	Enclosed: Yes
		Soil map as Annexure 3	Enclosed: Yes / No




## Annexure - I




**Fig: Mandal wise location map of the Prakasham district**

## Annexure – II



**Fig: Normal rainfall map of the Prakasham District**

### Annexure – III



**Fig: Soil map of the Prakasham District**

## 2.0 Strategies for weather related contingencies

### 2.1 Drought

#### 2.1.1 Rainfed situation

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 2 weeks (June 4 <sup>th</sup> week)	Rainfed Red soils	Redgram	No change	-	<b><u>Redgram varieties:</u></b> LRG-41, LRG-52, PRG-176, BSMR 736 <b><u>Castor hybrid:</u></b> PCH111, PCH 222 <b><u>Blackgram varieties:</u></b> TBG-104, GBG-1,PU-31, IPU 2-43 <b><u>Bengalgram:</u></b> JG-11,JAKI 9218,NBeG-49,3
		Redgram+ Castor (1:2)			
		Redgram + Foxtail millet (1:4)			
		Redgram + Bajra (1:2)			
		Castor + Bajra (1:2)			
		Castor			
		Fallow-FCV Tobacco (Rabi)			
		Blackgram			
	Rainfed Black soils	Cotton	Foxtail millet - Bengalgram	Sowing of foxtail millet (Var. Suryanadi) in the month of June preceding to bengalgram	
		Redgram			
		Fallow-Bengalgram (Rabi)			
		Fallow-Tobacco (FCV)(Rabi)			
		Blackgram	No change	--	

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
Early season drought (delayed onset)			Change in crop/cropping system	Agonomic measures	Remarks on Implementation
Delay by 4 weeks (July 2 <sup>nd</sup> week)	Rainfed Red soils	Redgram			
		Redgram+ Castor (1:2)			
		Redgram + Foxtail millet (1:4)			
		Redgram + Bajra (1:2)			

		Castor + Bajra (1:2)	No change	No change		
		Castor				
		Fallow-FCV Tobacco (Rabi)				
		Blackgram				
	Rainfed Black soils	Cotton				Sowing of foxtail millet (Var. Suryanadi) in the month of June and Use recommended seed rate to maintain optimum plant population
		Redgram				
		Fallow-Bengalgram ( <i>Rabi</i> )	Foxtail millet - Bengalgram			
		Fallow-Tobacco (FCV)(Rabi)	Foxtail millet - Bengalgram			
	Blackgram	No change				

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 6 weeks (July 4 <sup>th</sup> week)	Light soils-Rainfed	Redgram	No change	Reduce redgram row spacing from 120 cm to 90 cm	
		Redgram+ Castor (1:2)			
		Castor + Bajra (1:2)			
		Castor	No change	Reduce castor spacing from 90X60 cm to 90X45 cm	
		Castor + Bajra (1:2)			
		Fallow - FCV Tobacco (Rabi)			
		Blackgram	No change	--	
	Heavy soils-Rainfed	Cotton	No change	-	
		Fallow-Bengalgram (Rabi)	Foxtail millet - Bengalgram	Sowing of foxtail millet (Var. Suryanadi) in the month of June and Use recommended seed rate to maintain optimum plant population	
		Fallow-Tobacco (FCV)(Rabi)	Foxtail millet - Bengalgram		

		Blackgram	No change	Use recommended seed rate to maintain optimum plant population	
--	--	-----------	-----------	--	--

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 8 weeks (August 2 <sup>nd</sup> week)	Light soils-Rainfed	Redgram	No change	Reduce row spacing to 90 cm select LRG 52	
		Redgram+ Castor (1:2)	No change	-	
		Castor + Bajra (1:2)	No change	-	
		Castor	No change	Reduce spacing from 90X60 cm to 90X45 cm	
		Fallow - FCV tobacco (Rabi)	Foxtail millet - Bengalgram	Sowing of foxtail millet (Var. Suryanadi) in the month of June	
		Blackgram	Horsegram (or) Jowar (or) Foxtail millet (or) Bajra	Use recommended seed rate to maintain optimum plant population with short duration varieties	
	Heavy soils-Rainfed	Cotton	No change	Adopt closer spacing of 90X45 cm	
		Fallow - Bengalgram (Rabi)	Foxtail millet - Bengalgram	Sowing of foxtail millet in the month of August with short duration variety ( SIA 3222)	
		Fallow -Tobacco (Rabi)	Foxtail millet - Bengalgram	Sowing of foxtail millet in the month of August with short duration variety ( SIA 3222)	
		Blackgram	Chilli or Redgram or Foxtail millet	Redgram with LRG 52	

Condition			Suggested Contingency measures		
Early season drought (Normal onset)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
<b>Normal onset followed by 15-20 days dry spell after sowing leading to poor germination/crop stand etc.</b>	Heavy soils - Rainfed	Cotton	Gap filling	<ul style="list-style-type: none"> <li>When the crop is 2 weeks old take up inter cultivation to conserve moisture</li> <li>Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> <li>Formation of dead furrows</li> <li>If possible, give life saving irrigation</li> </ul>	
		Redgram	--	<ul style="list-style-type: none"> <li>When the crop is 2 weeks old take up inter cultivation to conserve moisture</li> <li>Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> <li>If possible, give life saving irrigation</li> </ul>	
		Blackgram	--	<ul style="list-style-type: none"> <li>When the crop is 2 weeks old take up inter cultivation to conserve moisture</li> <li>Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> <li>If possible, give life saving irrigation</li> </ul>	
		Bengalgram	--	<ul style="list-style-type: none"> <li>When the crop is 2 weeks old take up inter cultivation to conserve moisture</li> <li>Spray 2 % urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> </ul>	
	Light soils - Rainfed	Redgram (sole crop)		<ul style="list-style-type: none"> <li>Inter cultivation to be done after 2 weeks of sowing to conserve soil moisture</li> <li>Formation of dead furrows</li> <li>Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> </ul>	

		Redgram+ castor		<ul style="list-style-type: none"><li>• Inter cultivation to be done after 2 weeks of sowing to conserve soil moisture</li><li>• Formation of dead furrows</li><li>• Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li></ul>	
		Castor			
		Castor + Bajra			
		Blackgram			

Condition			Suggested Contingency measures		
Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
At vegetative stage	Heavy soils-rainfed	Cotton	Spray 2 % urea solution (or) 1 % water soluble fertilizers like KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Inter cultivate periodically (7-10 days interval) to conserve soil moisture</li> <li>• Protective irrigation</li> <li>• Formation of dead furrows</li> </ul>	
		Redgram	Spray 2% urea solution (or) 1% water soluble fertilizers like KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Frequent inter cultivation to conserve moisture</li> <li>• Protective irrigation</li> <li>• Formation of dead furrows</li> </ul>	



		Blackgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like KNO <sub>3</sub>	<ul style="list-style-type: none"><li>• Frequent inter cultivation to conserve moisture</li><li>• Protective irrigation</li><li>• Formation of dead furrows</li></ul>	
		Bengalgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like KNO <sub>3</sub>	<ul style="list-style-type: none"><li>• Protective irrigation</li></ul>	
	Light soils - rainfed	Redgram (sole crop)	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	<ul style="list-style-type: none"><li>• Frequent inter cultivation to conserve moisture</li><li>• Protective irrigation</li></ul> <p>Formation of dead furrows</p>	
		Redgram + castor inter crop	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>		
		Castor	<ul style="list-style-type: none"><li>• Spray 2% urea solution (or) 1% water soluble fertilizers KNO<sub>3</sub></li><li>• Adopt nipping to allow main spike to develop</li></ul>		
		Blackgram	<ul style="list-style-type: none"><li>• Spray 2% urea solution (or) 1% water soluble fertilizers KNO<sub>3</sub></li></ul>		
		Castor + bajra(1:2)	<ul style="list-style-type: none"><li>• Harvest intercrops as fodders as chances of grain yield are poor</li><li>• Supplement the nutrients to the main crop through foliar spray(Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub>)</li></ul>	Inter cultivate periodically (7-10 days interval) to conserve soil moisture	
		Redgram+ Castor (1:2)	<ul style="list-style-type: none"><li>• Adopt nipping to allow main spike to develop</li><li>• Supplement the nutrients to the main crop through foliar spray(Spray 2%</li></ul>		

			urea solution (or) 1% water soluble fertilizers like KNO <sub>3</sub> )		
--	--	--	---	--	--

Condition			Suggested Contingency measures		
Mid season drought (long dry spell)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
At reproductive stage	Heavy soils-rainfed	Cotton	Spray 2% urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Inter cultivate periodically (7-10 days interval) to conserve soil moisture</li> <li>• Protective irrigation</li> <li>• Formation of dead furrows</li> </ul>	
		Redgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Frequent inter cultivation to conserve moisture</li> <li>• Protective irrigation</li> <li>• Formation of dead furrows</li> </ul>	
		Blackgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Frequent inter cultivation to conserve moisture</li> <li>• Protective irrigation</li> <li>• Formation of dead furrows</li> </ul>	
		Bengalgram	Spray 2% urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Protective irrigation</li> </ul>	
	Light soils -rainfed	Redgram (sole crop)	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	<ul style="list-style-type: none"> <li>• Frequent inter cultivation to conserve moisture</li> <li>• Protective irrigation</li> </ul>	
		Redgram + castor inter crop	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	Formation of dead furrows	

		Castor	<ul style="list-style-type: none"> <li>• Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> <li>• Adopt nipping to allow main spike to develop</li> </ul>		
		Blackgram	<ul style="list-style-type: none"> <li>• Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub></li> </ul>		
		Castor + Bajra(1:2)	<ul style="list-style-type: none"> <li>• Harvest intercrops as fodders as chances of grain yield are poor</li> <li>• Supplement the nutrients to the main crop through foliar spray (Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub>)</li> </ul>	Inter cultivate periodically (7-10 days interval) to conserve soil moisture	
		Redgram+ Castor (1:2)	<ul style="list-style-type: none"> <li>• Adopt nipping to allow main spike to develop</li> <li>• Supplement the nutrients to the main crop through foliar spray(Spray 2% urea solution (or) 1% water soluble fertilizers like 19-19-19 (or) KNO<sub>3</sub>)</li> </ul>		
<b>Condition</b>			<b>Suggested Contingency measures</b>		
<b>Terminal drought</b>	<b>Major Farming situation</b>	<b>Normal Crop/cropping system</b>	<b>Crop management</b>	<b>Rabi Crop planning</b>	<b>Remarks on Implementation</b>
	Heavy soils-rainfed	Cotton	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	Rabi redgram/ Bengalgram / Safflower/ Coriander/Blackgram/Cowpea	
		Redgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	Rabi redgram/ Bengalgram / Safflower/ Coriander	

		Blackgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	Cowpea/ Bengalgram/ Safflower/ Sesamum/Coriander	
		Bengalgram	Spray 2 % urea solution (or) 1 % water soluble fertilizers like 19-19-19 (or) KNO <sub>3</sub>	--	
	light soils-Rainfed	Redgram (Sole crop)	Spray urea - 2 % or KNO <sub>3</sub> 1% or other water soluble fertilizers 1 % to supplement nutrition Selection of varieties with less duration if terminal drought is a common phenomenon	FCV-Tobacco / Cowpea/Foxtail millet/Greengram/Maghi jowar	
		Redgram+ Castor			
		Castor	1. Nipping of auxiliary buds to allow the main spike to mature 2. Foliar spray of urea 2 % or 1% KNO <sub>3</sub>	FCV-Tobacco/Cowpea/Foxtail millet/Greengram/Maghi jowar	
		Blackgram	Spray urea - 2 % or KNO <sub>3</sub> 1% or other water soluble fertilizers 1 % to supplement nutrition	FCV-Tobacco / Cowpea/Foxtail millet/Greengram/Maghi jowar	

### 2.1.2 Irrigated situation

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agonomic measures	Remarks on Implementation
Delayed release of water in canals due to low rainfall	Black soils – Canal irrigated (KWD)	Rice	Green manure-Rice-Greengram/Fodder	1.Green manure preceding to kharif rice Adopt preventive control measures for diseases like Blast in rice 2. Greengram i.e LGG - 460, WGG-42 can be grown in rice fallows under late seasonal conditions	-

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
	Red Soils/Black Soils – Canal irrigated (NSP Command)	Greengram/Green manure – Rice-Fodder	Greengram/Green manure – Rice - blackgram	1 Adopt medium duration varieties like NLR 34449, NLR 3041, NLR 145, JGL 384, MTU 1153, MTU 1156, NLR 4001, CR 1009 etc. which are resistant to blast and suitable for mid kharif season 2. If BPT 5204 is grown, timely plant protection is crucial	

Condition			Suggested Contingency measures		
Limited release of water in canals due to low rainfall	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
	Black soils – Canal irrigated (KWD)	Rice-Fodder	Rice (Semidry rice)-Greengram	<b>Rice</b> –1. Adopt alternate wetting and drying upto Primordial initiation then maintain upto a depth of 3 – 5 cm from Primordial initiation to maturity 2. Take up effective weed control measures either mechanically or through herbicides as the problem of weeds is more under alternate wetting and drying method of irrigation (specify herbicides and its concentration) <b>Rice fallows</b> 1. Crops like Greengram, Jowar etc. which require less water than Maize shall be grown 3. Short duration varieties of crops shall be selected 4. Protected irrigation can be given by using PVC/Metallic	<b>Rice</b> - weed management should be done using selective herbicides. <b>Rice fallows</b> – 1. Availability of seed of short duration varieties shall be ensured 2. Facilities like micro irrigation systems – Sprinkler and Drip can be extended to the farmers

				pipes instead of running water.	
--	--	--	--	---------------------------------	--

Condition	Suggested Contingency measures				
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Non release of water in canals under delayed onset of monsoon in catchment	Black soils/Red soils – Canal irrigation (NSP)	Rice	Bengalgram / redgram / Blackgram / Foxtail millet	Use recommended seed rate to maintain optimum plant population	
	Black soils – Canal irrigated (KWD)	Rice	Bengalgram / redgram / Blackgram / Foxtail millet	Use recommended seed rate to maintain optimum plant population	

Condition	Suggested Contingency measures				
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Lack of inflows into tanks due to insufficient /delayed onset of monsoon	Heavy soils – irrigated-tankfed	Redgram/Rice	Bengalgram /cotton/ redgram / Blackgram / Foxtail millet/ Castor/Sunflower	1. Use recommended seed rate to maintain optimum plant population 2. Foliar spray of nutrients 2% Urea or 1% KNO <sub>3</sub>	--
	Light soils- irrigated-tankfed/wells and bore wells	Summer Cotton-paddy	Redgram / Blackgram / Foxtail millet/Sunflower	1. Timely sowing is advantageous 2. Irrigation at critical stages through Micro irrigation systems	
		Cotton	Maghi Jowar/ Foxtail millet/ Cowpea/ Blackgram/	Use recommended seed rate to maintain optimum plant population	
		Bengalgram	Bengalgram	1. Timely sowing is advised 2. Irrigation at critical stages through sprinkler irrigation systems	

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agonomic measures	Remarks on Implementation
Insufficient groundwater recharge due to low rainfall	Bore well irrigated red soils and black soils	Groundnut	Blackgram/ Greengram/ /Sesamum/ Cotton/	1. Timely sowing is advised 2. Irrigation at critical stages through Micro irrigation systems 3. Irrigation at critical stages may be followed instead of intensive irrigations	
		Maize	Bengalgram/Cowpea/Jowar (Maghi)		
		Rice			
		Chilli	Blackgram/ Greengram/ /Sesamum/ Bengalgram		

## 2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigated situations)

Condition - Continuous high rainfall in a short span leading to water logging				
Crop	Suggested contingency measure			
	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Rice	1. Drain the excess water as early as possible 2. Apply 20 kg N + 10 kg K /acre after draining excess water 3. Take up gap filling either with available nursery or by splitting the tillers from the surviving hills 4. Take up proper weed control Measures 5. Timely plant protection measures in anticipation of pest & disease out breaks	1. Drain the excess water as early as possible 2. Apply 20 kg N + 10 kg K /acre after draining excess water 3. Take up suitable plant protection Measures in anticipation of pest & disease out breaks	1. Drain the excess water as early as possible 2. Take up suitable plant protection measures in anticipation of pest & disease out breaks	1. Drain out water and spread sheaves loosely in field or field bunds where there is no water stagnation 2. Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds 3. Thresh after drying the sheaves properly 4. Ensure proper grain moisture before storing
Cotton	1. Drain the excess water as early as possible in black soils 2. Apply 20 kg N + 15 kg K /ha after draining excess water 3. Take up inter cultivation at optimum soil moisture	1. Drain the excess water as early as possible 2. Apply 20 kg N + 15 kg K /ha after draining excess water 3. To spray 1 % KNO <sub>3</sub> or 2% Urea to support nutrition 4. Spray fungicides like Copper oxy	1. Drain the excess water as early as possible 2. To spray 1 % KNO <sub>3</sub> or 2% Urea to support nutrition 3. Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or	Dry the produce properly before baling and sending to market

	<p>condition to loosen and aerate the soil and to control weeds</p> <ol style="list-style-type: none"> <li>To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</li> <li>Take up timely control measures against sucking pests</li> </ol>	<p>chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals to control Bacterial leaf blight, wilt alternaria leaf spot and grey mildew</p> <ol style="list-style-type: none"> <li>Take up timely control measures against sucking pests and bollworms.</li> </ol>	<p>Mancozeb 0.25% against boll rot</p> <ol style="list-style-type: none"> <li>Take up timely control measures against bollworms and whitefly</li> </ol>	
Redgram	<ol style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Apply 20 kg N + 10 kg K /acre after draining excess water</li> <li>Take up inter cultivation at optimum soil moisture status to loosen and aerate the soil and to control weeds</li> <li>To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</li> </ol>	<ol style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>Take up timely control measures against possible outbreak of pod borer complex, maruca , Helicovera etc.</li> </ol>	<ol style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Allow the crop to dry completely before harvesting</li> </ol>	<ol style="list-style-type: none"> <li>Spread the bundles drenched in rain on field bunds or drying floors to quicken the drying</li> <li>Thresh the bundles after they are dried properly</li> <li>Dry the grain to proper moisture per cent before bagging and storing to prevent deterioration in quality during storage</li> </ol>
Castor	<ol style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Apply 4-5 kg N /acre after draining excess water</li> <li>To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>Take up timely control measures for semilooper</li> </ol>	<ol style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Apply 4-5 kg N /acre after draining excess water</li> <li>To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.2 % for Botrytis grey rot control</li> <li>Take up timely control measures</li> </ol>	<ol style="list-style-type: none"> <li>Drain the excess water as early as possible</li> <li>Allow the crop to dry completely before harvesting</li> </ol>	



		against <i>Spodoptera</i> and capsule borer		
Maize	<ol style="list-style-type: none"> <li>1. Drain the excess water as early as possible</li> <li>2. Apply 20 kg N + 10 kg K /acre after draining excess water</li> <li>3. Take up inter cultivation and at optimum soil moisture condition to loosen and aerate the soil and to control weeds</li> <li>4. To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</li> <li>5. Take up timely control measures for Pink stem borer, sheath blight</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain the excess water as early as possible</li> <li>2. Apply 20 kg N + 10 kg K /acre after draining excess water</li> <li>3. To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</li> <li>4. Take up timely control measures for sheath blight and post flowering stalk rots</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain the excess water as early as possible</li> <li>2. Allow the crop to dry completely before harvesting</li> </ol>	<ol style="list-style-type: none"> <li>1. Harvest the cobs after the they are dried up properly. Dry the grain to optimum moisture condition before storing</li> </ol>
Bengalgram	<ol style="list-style-type: none"> <li>1. Drain the excess water as early as possible</li> <li>2. To spray KNO<sub>3</sub> 1 % to support nutrition</li> <li>4. Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25%</li> <li>5. Take up timely control measures against the out break of pests like <i>Helicoverpa</i> etc.</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain the excess water as early as possible</li> <li>2. To spray KNO<sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</li> <li>4. Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% against blight and wilt</li> <li>5. Take up timely control measures against the outbreak of pests like <i>Helicoverpa</i> etc.</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain the excess water as early as possible</li> <li>2. Allow the crop to dry completely before harvesting</li> </ol>	
Horticulture (Fruits)				
Orange & Batavian	<p>Drain the excess water as soon as possible.</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Foliar spray of micronutrient</p>	<p>Drain the excess water as soon as possible.</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Foliar spray of micronutrient</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature fruits in a clear sunny day.</p>	<p>Store the fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p>

	<p>mixture is also to be taken up.</p> <p>Sand casting around the tree trunks should be removed up to the collar region of the tree to prevent fungal infections.</p> <p>If the tree age is above eight years a booster dose of 500 g of Urea and 750 g MOP per tree should be applied.</p>	<p>mixture is also to be taken up.</p> <p>Sand casting around the tree trunks should be removed up to the collar region of the tree to prevent fungal infections.</p> <p>If the tree age is above eight years a booster dose of 500 g of Urea and 750 g MOP per tree should be applied.</p>	.	
Mango	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	Same as above	Same as above
Sapota	Same as above	Same as above	Same as above	Same as above
Lemon	<p>Drain the excess water as soon as possible.</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Foliar spray of micronutrient mixture is also to be taken up.</p> <p>Sand casting around the tree trunks should be removed up to the collar region of the tree to prevent fungal infections.</p> <p>If the tree age is above eight years a booster dose of 500 g of Urea and 750 g MOP per tree should be applied.</p>	<p>Drain the excess water as soon as possible.</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Foliar spray of micronutrient mixture is also to be taken up.</p> <p>Sand casting around the tree trunks should be removed up to the collar region of the tree to prevent fungal infections.</p> <p>If the tree age is above eight years a booster dose of 500 g of Urea and 750 g MOP per tree should be applied.</p>	.	
Papaya	<p>Drain out the excess water</p> <p>out break of any sucking past</p>	<p>Drain out the excess water</p> <p>Water logging near trunk should be prevented</p>	<p>Drain out the excess water</p> <p>Harvest the marketable fruits in a</p>	

	<p>should be controlled using systemic insecticides</p> <p>Water logging near trunk should be prevented</p>		<p>clear sunny day</p> <p>Water logging near trunk should be prevented</p> <p>Micronutrient deficiencies should be corrected by foliar sprays of Fe, Mg, Zn, Bo and Mn</p>	
Vegetables				
Chillies	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.</p> <p>In case of severe damage (considered as complete economical loss), and the contingency period is between June to August, sowing of best alternative crop must be taken up.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the matured fruits in a clear sunny day.</p>	<p>Dry the pods on concrete floor immediately after the appearance of sunlight (or).</p> <p>Use poly house solar driers for quick drying</p> <p>Grade the pods and market as soon as possible.</p> <p>Do not store such produce for long periods.</p>
Tomato	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 12 kg MOP + 30 kg Urea per acre as soon</p>	Same as above	Same as above	<p>Store the harvested fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p>

	<p>as possible.</p> <p>Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.</p> <p>In case of severe damage (considered as complete economical loss), and the contingency period is between June to August, sowing of best alternative crop must be taken up.</p>			
<b>Spices and plantation crops</b>				
Coriander	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the marketable umbels as soon as possible.</p>	<p>Dry the produce immediately</p> <p>Market the produce immediately after drying.</p>
<b>Condition - Heavy rainfall with high speed winds in a short span</b>				
Rice	Measures similar to above given for heavy rainfall situation as above	In addition to the above measures lift the lodged hills and tie them together to keep them erect	In addition to the above measures, lift the lodged plants and tie them together keep erect	In addition to the above measures, for water lagged take up measures to minimize blowing away of produce due to high velocity winds.
Cotton	<p>In addition to the measures for removing excess water,</p> <p>Lift the fallen plants if any and firm up the soil around the base of the stem</p>	<p>Lift the fallen plants if any and firm up the soil around the base of the stem</p> <p>Bacterial leaf blight: Spray plantomycin 16g per acre</p>	Similar measures as in water lagged situation. Additional by pick the net cotton at the earliest	Dry the produce under sun before sending to market

Redgram	<p>Lift the lodged plants if any and firm up the soil around the base of the stem</p> <p>Apply 4-5 kg N /acre after draining excess water</p>	<p>Lift the lodged plants if any and firm up the soil around the base of the stem</p> <p>Takeup timely pest control measures for pod borers and wilt</p>	Harvest the pods from uprooted plants as soon as the field condition permits and transport to drying floor	Dry the produce under sun before thrashing and sending to market.
Castor	<p>1. Drain out the excess water from the field as early as possible</p> <p>2. Apply 4-5 kg N/acre after draining excess water</p> <p>3. To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</p> <p>4. Take up proper weed control measures</p> <p>5. Take up timely plant protection measures for possible pest and disease out breaks</p>	<p>1. Drain out the excess water from the field as early as possible</p> <p>2. Apply 4-5 kg N /acre after draining excess water</p> <p>3. To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</p> <p>4. Take up timely plant protection measures for possible pest and disease out breaks</p>	<p>1. Drain out the excess water from the field as early as possible</p> <p>2. Harvest the crop as soon as the field condition permits</p>	1. Dry the produce under sun before sending to market
Maize	<p>Drain out the excess water from the field as early as possible</p> <p>Earthing-up for better anchorage</p>	Drain out the excess water from the field as early as possible	<p>Drain out the excess water from the field as early as possible</p> <p>Allow the crop to dry completely before harvesting</p>	Harvest the cobs after they are dried up properly. Dry the grain to optimum moisture condition before storing
Horticulture				
Orange & Batavian	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> followed by Ferrous Sulphate 0.5% + Citric Acid 0.1 % solution 2-3</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> followed by Ferrous Sulphate 0.5%</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Harvest the matured fruits in a clear day by using improved</p>	<p>Grade the damaged or infected fruits.</p> <p>Store the graded fruits in well-ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p> <p>The fallen under sized</p>

	<p>times.</p> <p>Topdressing of booster dose of 40 kg MOP + 50 kg Urea along with 250 kg of Neem Cake per acre as soon as possible.</p>	<p>+ Citric Acid 0.1 % solution 2-3 times.</p> <p>Topdressing of booster dose of 40 kg MOP + 50 kg Urea along with 250 kg of Neem Cake per acre as soon as possible.</p>	harvesters	fruits may be utilized for processing immediately
Mango	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	Same as above	Same as above
Guava	<p>Provide support to the young plants</p> <p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Harvest the mature fruits as soon as possible.</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	Same as above
Lemon	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p>	<p>Wind damaged branches should be pruned using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p>	Same as above

	<p>Spray Urea 2% or 1% KNO<sub>3</sub> followed by Ferrous Sulphate 0.5% + Citric Acid 0.1 % solution 2-3 times.</p> <p>Topdressing of booster dose of 40 kg MOP + 50 kg Urea along with 250 kg of Neem Cake per acre as soon as possible.</p>	<p>Spray Urea 2% or 1% KNO<sub>3</sub> followed by Ferrous Sulphate 0.5% + Citric Acid 0.1 % solution 2-3 times.</p> <p>Topdressing of booster dose of 40 kg MOP + 50 kg Urea along with 250 kg of Neem Cake per acre as soon as possible.</p>	Harvest the matured fruits in a clear day	
Papaya	<p>Uprooted plants may be lifted and earthed up</p> <p>Gap filling\Replanting may be done based on extent of damage</p> <p>Stake the plants if necessary</p>	Staking may be provided for heavy bearing plants	<p>Same as above and</p> <p>Staking may be provided for heavy bearing plants</p> <p>Dropped fruits should be collected from garden</p>	<p>Drain the excess water as soon as possible.</p> <p>Grade the damaged or infected fruits.</p> <p>Store the graded fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p> <p>The fallen under sized fruits may be utilized for processing immediately.</p>
Vegetables				
Chillies	<p>Uprooted plants may be lifted and earthed up</p> <p>Gap filling must be done immediately</p> <p>If damage is more, go for replanting</p> <p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or KNO<sub>3</sub> 1%</p>	<p>Uprooted plants may be lifted and earthed up</p> <p>Gap filling must be done immediately</p> <p>If damage is more ,go for replanting</p> <p>Drain the excess water as soon as possible</p>	<p>Uprooted plants may be lifted and earthed up</p> <p>Drain the excess water as soon as possible</p> <p>Harvest the matured fruits in a clear sunny day.</p>	<p>Dry the pods on elevated concrete floor\polythene sheet immediately after the appearance of sunlight (or).</p> <p>Use poly house solar driers for quick drying</p> <p>Dry the chillies till it produces rattling sound</p>

	<p>solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Intercultivate the soil with gorru and guntaka for better aeration</p>	<p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p>		<p>(10-11% moisture)</p> <p>Grade the pods and market as soon as possible.</p> <p>Do not store such produce for long periods. .</p>
Tomato	<p>Uprooted plants may be lifted and earthed up</p> <p>Gap filling must be done immediately</p> <p>If damage is more, go for replanting</p> <p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution once.</p>	<p>Uprooted plants may be lifted and earthed up</p> <p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution once.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the marketable fruits in a clear sunny day.</p>	<p>Store the harvested fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p>
Spices and plantation crops				
Coriander	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> solution 2-3 times.</p>	<p>Drain the excess water as soon as possible</p> <p>Harvest the marketable umbels as soon as possible.</p>	<p>Dry the produce immediately</p> <p>Market the produce immediately after drying.</p>
<b>Condition - Outbreak of pests and diseases due to unseasonal rains</b>				
Rice	Stem borer, Leaf folder	Stem borer, Leaf folder , stem rot , BLB,BLS, BPH incidence is more	Blast, stem rot and Panicle mite	
Cotton	Jassids, Aphids, Thrips	Jassids , Thrips, Pink boll worm	Pink boll worm ,Spodoptera, <i>Helicoverpa</i>	
Redgram	Wilt	SMD, Wilt	Maruca and Pod borer Helicoverpa Wilt	
Castor	---	Leaf spots	Hairy caterpillar and Botrytis	



			grey rot	
Sunflower	Jassids, aphids	Green caterpillar, Aphids, Leaf blight and bud necrosis	Bud necrosis, <i>Helicoverpa</i>	
Bengalgram	<i>Spodoptera exigua</i> , Wilt , Blight	<i>Helicoverpa</i> Wilt	<i>Helicoverpa</i> , wilt	
Blackgram	Thrips, YMV	YMV, Thrips , Bud necrosis	YMV	
Horticulture				
Orange, Batavian, Lemon	-	Bacterial leaf spot	Orange, Batavian, Lemon	
Mango	Hoppers, Thrips	Anthrachnose	Mango	
Sapota	Whitefly, mealy bug, fruit fly	Anthrachnose, wilt	Sapota	
papaya		Ring spot virus	papaya	
Chillies	Thrips, mites, Spodoptera and <i>Helicoverpa</i>	Die back and fruit rot, Bacterial leaf spot, viruses	Chillies	
Tomato	<i>Helicoverpa</i>	Blight, wilt, virus	Tomato	

### 2.3 Floods

Condition	Transient water logging/ partial inundation			
	Suggested contingency measure			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Rice	1. To drain out the excess water at the earliest 2. Apply booster dose of 0.5 kg N/40 sq. m 3. Spray micronutrients like Zn, Fe two to three times at 4 -5 days interval 4. Take up proper weed control measures	1. To drain out the excess water at the earliest 2. Take up gap filling either with available nursery or by splitting the tillers from the surviving hills 3. Apply booster dose of 20 kg N/Acre 4. Spray ZnSO <sub>4</sub> 0.2 % if it is less than 45 days after transplanting 5. Take up need based plant protection measures	1. To drain out the excess water at the earliest 2. Take up need based plant protection measures	1. Drain out water spread sheaves loosely in field or field bunds where there is no water stagnation 2. Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds 3. Thresh after drying the sheaves properly 4. Ensure proper grain moisture before storing

Cotton	<ol style="list-style-type: none"> <li>1.To drain out the excess water at the earliest by farming drainage channels if there is a gradient and if not by using motors</li> <li>2. Take up the gap filling at the earliest</li> <li>3.Inter cultivate at optimum field moisture condition</li> <li>4.Apply 20 kg N + 15 kg K /ha after draining excess water</li> <li>5.To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>6.Take up plant protection measures against possible pests and disease incidence</li> <li>7.Select short duration hybrids</li> <li>8.Adopt closer spacing of 90 X 45 or 90 X 30 cm</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest by farming drainage channels if there is a gradient and if not by using motors Inter cultivate at optimum field moisture condition</li> <li>2. Apply 20 kg N + 15kg K /ha after draining excess water</li> <li>3. To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>4. Spray of micronutrients two times at 7-10 days interval</li> <li>5. Take up plant protection measures against possible pests and disease incidence</li> </ol>	<ol style="list-style-type: none"> <li>1.To drain out the excess water at the earliest by farming drainage channels if there is a gradient and if not by using motors 5</li> <li>2.To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>3.Take up plant protection measures against possible pests and disease incidence</li> </ol>	Kapas picking should be done carefully to prevent admixtures with waste plant material
Redgram	<ol style="list-style-type: none"> <li>1.To drain out the excess water at the earliest</li> <li>2. Take up the gap filling at the earliest</li> <li>3.Inter cultivate at optimum field moisture condition</li> <li>4.Apply 4-5 kg N/acre after draining excess water</li> </ol>	<ol style="list-style-type: none"> <li>1.To drain out the excess water at the earliest</li> <li>2.Take up the gap filling at the earliest</li> <li>3.Inter cultivate at optimum field moisture condition</li> <li>4.Apply 4-5 kg N/acre after draining excess water</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>3. Take up plant protection measures against possible pests and disease incidence</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Harvest the crop when the field condition permits</li> <li>3. Drying of bundles should be done on elevated places like filed bunds or drying floors</li> </ol>
Bengalgram	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Take up the gap filling at the earliest</li> <li>3.Take up weed control either</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Takeup weed control either mechanically or through</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Apply 4-5 kg N/acre</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain out the excess water at the earliest</li> <li>2. Harvest the crop after the fields are dried up</li> </ol>

	mechanically or through weedicides 4. Apply 4-5 kg N/acre after draining excess water 5. Take up plant protection measures against possible pests and disease incidence	weedicides 3. Apply 4-5 kg N/acre after draining excess water 4. To spray 1 % KNO <sub>3</sub> or 2% Urea to support nutrition 5. Take up plant protection measures against possible pests and disease incidence	after draining excess water 3. To spray 1 % KNO <sub>3</sub> or 2% Urea to support nutrition 4. Take up plant protection measures against possible pests and disease incidence	
Sunflower	1. To drain out the excess water at the earliest 2. Takeup the gap filling at the earliest 3. Inter cultivate at optimum field moisture condition 4. Apply 4-5 kg N/acre after draining excess water	1. To drain out the excess water at the earliest 2. Takeup the gap filling at the earliest 3. Inter cultivate at optimum field moisture condition 4. Apply 4-5 kg N/acre after draining excess water	1. To drain out the excess water at the earliest 2. To spray 1% KNO <sub>3</sub> or 2% Urea to support nutrition 3. Take up plant protection measures against possible pests and disease incidence	1. To drain out the excess water at the earliest 2. Harvest the crop when the field condition permits 3. Drying of bundles should be done on elevated places like filed bunds or drying floors
<b>Condition - Continuous submergence for more than 2 days</b>				
	<b>Suggested contingency measure</b>			
Rice	1. Top dressing with 0.5 kg N/40 sq.m immediately after recede of flood water 2. Spray of ZnSO <sub>4</sub> , FeSO <sub>4</sub> to correct micronutrient deficiencies 3. Weed control through mechanical or Chemical measures	1. To drain out the excess water at the earliest 2. Take up gap filling either with available nursery or by splitting the tillers from the surviving hills if the gaps are < 30% if more go for replanting 3. Apply 20 kg N + 10 kg K /acre after draining excess water 4. Proper weed control measures to be taken up 4. Timely plant protection measures for pest and disease out break	1. To drain out the excess water at the earliest 2. Take up need based plant protection measures	1. Drain out water spread sheaves loosely in field or field bunds where there is no water stagnation 2. Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds 3. Thresh after drying the sheaves properly 4. Ensure proper grain moisture before storing

Cotton	<ol style="list-style-type: none"> <li>1. Mortality is most likely hence re sowing to be taken up</li> <li>2. Select short duration hybrids</li> <li>3. Adopt closer spacing of 90X45 &amp; 90X30 cm</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Apply 20 kg N + 10 kg K /acre after draining excess water</li> <li>3. Spray micronutrient mixture for 2 to 3 times at an interval of 7-10 days</li> <li>4. To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>5. Inter cultivate to smother weeds and to loosen and aerate the soil</li> <li>6. Need based plant protection measures to be taken up</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Spray micronutrient mixture for 2 to 3 times at an interval of 7-10 days</li> <li>3. To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>4. Need based plant protection measures to be taken up</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain out the water as early as possible</li> <li>2. To spray 1% KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>3. Kapas picking should be done carefully to avoid admixtures and plant waste</li> </ol>
Redgram	<ol style="list-style-type: none"> <li>1. Take up gap filling if the gaps are &lt; 30 % and if more take up re sowing</li> <li>2. After gap filling take up inter cultivation to smother the weeds and to aerate the soil</li> <li>3. Apply 20 kg N + 10 kg K /acre after draining excess water</li> </ol>	<ol style="list-style-type: none"> <li>1. After gap filling take up inter cultivation to smother the weeds and to aerate the soil</li> <li>2. Apply 20 kg N + 10 kg K /acre after draining excess water</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain out excess water form the field</li> <li>2. Apply 20 kg N + 10 kg K /acre after draining excess water</li> <li>3. Need based plant protection measures to be taken up</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain out excess water as early as possible</li> </ol>
Bengalgram	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Takeup gap filling if the gaps are &lt; 30 % and if more take up resowing</li> <li>3. Apply 4-5 kg N /acre after draining excess water</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. Apply 4-5 kg N/acre after draining excess water</li> <li>3. To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>4. Proper weed control measures to be taken up</li> <li>5. Need based plant protection measures to be taken up</li> </ol>	<ol style="list-style-type: none"> <li>1. To drain out the excess water at the earliest</li> <li>2. To spray 1 % KNO<sub>3</sub> or 2% Urea to support nutrition</li> <li>3. Need based plant protection measures to be taken up</li> </ol>	<ol style="list-style-type: none"> <li>1. Drain out the excess water at the earliest</li> </ol>

Castor	1. To drain out the excess water at the earliest 2. Re sow the crop if mortality is > 15 % 3. Apply 20 kg N + 10 kg K /acre after draining excess water	1. To drain out the excess water at the earliest 2. Apply 20 kg N + 10 kg K /acre after draining excess water 3. Inter cultivate to smother weeds and to loosen and aerate the soil 4. To spray 1 % KNO <sub>3</sub> or 2% Urea to support nutrition 5. Need based plant protection measures to be taken up	1. To drain out the excess water at the earliest 2. Apply 20 kg N + 10 kg K /acre after draining excess water 3. To spray 1 % KNO <sub>3</sub> or 2% Urea to support nutrition 4. Need based plant protection measures to be taken up	Drain out the excess water at the earliest
Sunflower	1. Mortality is most likely hence re sowing to be taken up 2. Select short duration hybrids 3. Adopt closer spacing of 45 X 30 cm	1. To drain out the excess water at the earliest 2. Apply 20 kg N + 10 kg K /acre after draining excess water 3. Spray micronutrient mixture for 2 to 3 times at an interval of 7-10 days 4. To spray KNO <sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition 5. Inter cultivate to smother weeds and to loosen and aerate the soil 6. Need based plant protection measures to be taken up	1. To drain out the excess water at the earliest 2. Spray micronutrient mixture for 2 to 3 times at an interval of 7-10 days 3. Spray KNO <sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition 4. Need based plant protection measures to be taken up	1. Drain out the water as early as possible 2. To spray KNO <sub>3</sub> 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition
Horticulture				
Orange & Batavian, Mango, Guava, Lemon, Papaya	Drain the excess water as soon as possible.  Spray 1% KNO <sub>3</sub> or Urea 2% solution 2-3 times.	Drain the excess water as soon as possible.  Spray 1% KNO <sub>3</sub> or Urea	Drain the excess water as soon as possible.  Spray 1% KNO <sub>3</sub> or Urea	Drain the excess water as soon as possible.  Harvest the mature fruits as

		<p>2% solution 2-3 times.</p> <p>Foliar spray of micronutrient mixture is also to be taken up.</p> <p>Sand casting around the tree trunks should be removed up to the collar region of the tree to prevent fungal infections.</p> <p>If the tree age is above eight years a booster dose of 500 g of Urea and 750 g MOP per tree should be applied.</p>	<p>2% solution 2-3 times.</p> <p>Foliar spray of micronutrient mixture is also to be taken up.</p> <p>Sand casting around the tree trunks should be removed up to the collar region of the tree to prevent fungal infections.</p> <p>If the tree age is above eight years a booster dose of 500 g of Urea and 750 g MOP per tree should be applied.</p>	<p>soon as possible.</p> <p>Store the fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p>
Chillies	Drain the excess water as soon as possible	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Gap filling may be taken up if the plants are two weeks old and sowing window is still available for the crop.</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.</p> <p>Spray planofix 1ml in 4.5 lit of water to prevent flower drop.</p>	<p>Drain the excess water as soon as possible.</p> <p>Dry the pods on concrete floor/ tarpaulins.</p> <p>Spray any drying oil after the pods are free from surface moisture for quick drying.</p> <p>Use poly house solar driers for quick drying</p> <p>Remove the pest and disease infected pods.</p> <p>Market the produce as soon as possible</p>
Tomato	Same as above	Drain the excess water as	Drain the excess water	Same as above

		<p>soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 10 kg MOP + 30 kg Urea per acre as soon as possible.</p>	<p>as soon as possible</p> <p>Spray Urea 2% solution once.</p>	
Spices and plantation crops				
Coriander		<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> solution 2-3 times</p>	<p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% or 1% KNO<sub>3</sub> solution 2-3 times</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the marketable umbels as soon as possible.</p> <p>Dry the produce immediately</p> <p>Market the produce immediately after drying.</p>

#### 2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone

Extreme event type	Suggested contingency measurer			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Cyclone				
Horticulture crops – Fruits				
Orange & Batavian	<p>Spray Carbendazim 1 g or COC 3g per litre to prevent spread of diseases</p> <p>If the damage is severe, go for resowing.</p>	<p>Tress fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have</p>	<p>Tress fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature fruits as soon as possible.</p> <p>Collect the fallen fruits</p>

		<p>to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>and sell immediately or go for preparation of processed products.</p> <p>If to store, store the produce in well ventilated place temporarily before it can be marketed.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>
Mango	If the damage is severe, go for resowing	<p>Trees fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>Tress fallen on ground may be lifted and earthed up</p> <p>.Manuring and plant protection measures have to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature fruits as soon as possible.</p> <p>Collect the fallen fruits and sell immediately or go for preparation of processed products.</p> <p>If to store, store the produce in well ventilated place temporarily before it can be marketed.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>
Sapota	Drain the excess water as	Wind damaged branches should be pruned using	Wind damaged branches should be pruned using	Wind damaged branches should be pruned



	<p>soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p> <p>Provide support to the young plants.</p>	<p>disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>using disinfected secateurs and cut ends must be smeared with Bordeaux paste</p> <p>Drain the excess water as soon as possible.</p> <p>Harvest the mature fruits as soon as possible.</p> <p>Store the fruits in well ventilated place temporarily before it can be marketed.</p> <p>Market the fruits as soon as possible.</p> <p>The unmarketable fruits may be utilized for processing</p>
Lemon	<p>If the damage is severe, go for resowing.</p>	<p>Tress fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>Tress fallen on ground may be lifted and earthed up</p> <p>Manuring and plant protection measures have to be taken up.</p> <p>Broken and damaged branches may be pruned and applied with Bordeaux paste</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature fruits as soon as possible.</p> <p>Collect the fallen fruits and sell immediately or go for preparation of processed products.</p> <p>If to store, store the produce in well ventilated place temporarily before it can be marketed.</p>

				Broken and damaged branches may be pruned and applied with Bordeaux paste
Papaya		Spray 1% KNO <sub>3</sub> or Urea 2% solution 2-3 times.	<p>Drain the excess water as soon as possible</p> <p>Spray 1% KNO<sub>3</sub> or Urea 2% solution 2-3 times.</p>	<p>Drain the excess water as soon as possible.</p> <p>Harvest the mature produce as soon as possible.</p> <p>Store the produce in well ventilated place temporarily before it can be marketed.</p> <p>Market the produce as soon as possible.</p> <p>Collect the fallen fruits and sell immediately or go for preparation of processed products.</p>
Horticulture crops vegetables				
Chillies	Grow nursery on raised beds.	<p>Uprooted plants may be lifted and earthed up</p> <p>Drain the excess water as soon as possible</p> <p>Gap filling must be done immediately</p> <p>If damage is more go for replanting Spray Urea 2% solution 2-3</p>	<p>Uprooted plants may be lifted and earthed up</p> <p>Drain the excess water as soon as possible</p> <p>Spray Urea 2% solution 2-3 times.</p> <p>Topdressing of booster dose of 15 kg MOP + 30 kg Urea per</p>	<p>Drain the excess water as soon as possible.</p> <p>Dry the pods on concrete floor/ tarpaulins immediately</p> <p>use poly house solar driers for quick drying</p>

		times.  Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.	acre as soon as possible.	.
Tomato	Grow nursery on raised beds.  If damage is more go for resowing	Uprooted plants may be lifted and earthed up  Drain the excess water as soon as possible  Gap filling must be done immediately  Spray Urea 2% solution 2-3 times.  Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.  If damage is more ,go for replanting	Uprooted plants may be lifted and earthed up  Drain the excess water as soon as possible  Spray Urea 2% solution 2-3 times.  Topdressing of booster dose of 15 kg MOP + 30 kg Urea per acre as soon as possible.  If damage is more , go for replanting	Drain the excess water as soon as possible.  Harvest the mature produce as soon as possible.  Store the produce in well ventilated place temporarily before it can be marketed.  Market the produce as soon as possible.
Spices and Plantation crops				
Coriander		Drain the excess water as soon as possible  Spray Urea 2% or 1% KNO <sub>3</sub> solution 2-3	Drain the excess water as soon as possible  Spray Urea 2% or 1% KNO <sub>3</sub> solution 2-3 times.	Drain the excess water as soon as possible.  Harvest the marketable umbels as soon as possible.  Dry the produce

		times.		<p>immediately</p> <p>Market the produce immediately after drying.</p> <p>Spray Dithane M-45/ Bavistin to prevent grey mould on the standing crop.</p>
--	--	--------	--	--

## 2.5 Contingent strategies for Livestock, Poultry & Fisheries

### General contingency measures

Before the event	During the event	After the event
<b>Feed and fodder availability</b>		
<p>1.Conserving fodder/crop residues/ forest grass by silage/hay making either by individual or on community basis</p> <p>2. Preparing complete diets and storing in strategic locations</p> <p>3. Organize procurement of dry fodders / feed ingredients from surplus areas</p> <p>4. Establish fodder banks and feed banks</p> <p>5. Livestock relief camps during floods/cyclones must be planned in the vicinity of relief camps for people</p> <p>6. Capacity building and preparedness</p>	<p>1.Organise relief camps 2.Supply silage / hay to farmers with productive stock on subsidized rates</p> <p>3.Segregate old, weak and unproductive stock and send for slaughter</p> <p>4. Supply mineral mixture to avoid deficiencies</p> <p>5. Dry fodder must be offered to the livestock in little quantities for number of times</p> <p>6.Concentrate feed or complete feed must be offered to only productive and young stock only</p>	<p>1. Capacity building to stake holders on drought /cyclone/flood mitigation in livestock sector</p> <p>2. Promote fodder cultivation.</p> <p>3. Flushing the stock to recoup</p> <p>4. Avoid soaked and mould infected feeds / fodders to livestock</p> <p>5. Replenish the feed and fodder banks</p> <p>6.Promote fodder preservation techniques like silage / hay making</p>
<b>Drinking water</b>		

<p>1.Construct drinking water tanks in herding places, village junctions and in relief camp locations</p> <p>2.Plan for sufficient number of tanks for water transportation</p> <p>3. Identify bore wells, which can sustain demand.</p> <p>4.Procure sufficient quantities of water Sanitizers</p>	<p>1.Regular supply of clean drinking water to all tanks 2.Cleaning the tanks in regular intervals</p> <p>3.Keep the livestock away from contaminated flood/cyclone/stagnated waters</p> <p>3.Add water sanitizers</p>	<p>1.Hand over the maintenance of the structures to panchayats</p> <p>2.Sensitize the farming community about importance of clean drinking water</p>
<b>Health and disease Management</b>		
<p>1.Procure and stock emergency medicines and vaccines for important endemic diseases of the area</p> <p>2. All the stock must be immunized for endemic diseases of the area</p> <p>3. Carry out deworming to all young stock</p> <p>4. Keep stock of bleaching powder and lime</p> <p>5.Carry out Butax spray for control of external parasites</p> <p>6.Identify the Clinical staff and trained paravets and indent for their services as per schedules</p> <p>7.Identify the volunteers who can serve in need of emergency</p>	<p>1.Keep close watch on the health of the stock</p> <p>2. Sick animals must be isolated and treated separately.</p> <p>3. Carry out deworming and spraying to all animals entering into relief camps</p> <p>4. Clean the animal houses regularly and apply disinfectants.</p> <p>5.Safe and hygienic disposal of dead animal carcasses</p> <p>6. Organize with community daily lifting of dung from relief camps</p>	<p>1.keep close surveillance on disease outbreak.</p> <p>2.Undertake the vaccination depending on need</p> <p>3.Keep the animal houses clean and spray disinfectants</p>

## 2.5.1 Detailed contingency strategies for Livestock, Poultry & Fisheries

	Suggested contingency measures		
	Before the event	During the event	After the event
<b>Drought</b>			
Feed and Fodder availability	<p>Some mandals of the district are chronically drought prone. It should have reserves of the following at any point of the year for mobilization to the needy areas (for feeding 5000 ACU (maintenance ration) for about 1-3 weeks period)</p> <p>Silage:20-50 t</p> <p>Urea molasses mineral bricks (UMMB):50-100 t</p> <p>Hay:100-250 t</p> <p>Concentrates: 20-50 t</p> <p>Minerals and vitamin supplements mixture:1-5 t</p> <p>Establishment of silvi-pastoral system in CPRs with <i>Stylosanthes hamata</i> and <i>Cenchrus ciliaris</i> as grass with <i>Leucaena leucocephala</i> as tree component (or suggest suitable similar system to your district)</p> <p>Top dressing of N in 2-3 split doses @ 20-25 kg N/ha in common property resources (CPRs) like temple lands, panchyat lands or private property resources (PPRs) like waste and degraded lands with the monsoon pattern for higher biomass production</p> <p>Promote cultivation of short duration fodder crops of sorghum/bajra/maize(UP chari, MP chari, HC-136, HD-2, GAIN BAJRA, L-74, K-677, Ananad/African Tall, Kisan composite, Moti, Manjari, B1-7 and also sunhemp</p> <p>Chopping of fodder should be made as mandatory in every village through supply and establishment of</p>	<p>Harvest and use biomass of dried up crops (Rice, Maize, Bajra, Horse gram, Groundnut, black gram, sun hemp) material as fodder.</p> <p>Harvest the tree fodder (Neem, Subabul, Acasia, Pipal etc) and unconventional feeds resources available and use as fodder for livestock (LS).</p> <p>Available feed and fodder should be cut from CPRs and stall fed in order to reduce the energy requirements of the animals</p> <p>UMMB, hay, concentrates and vitamin &amp; mineral mixture should be transported to the needy areas from the reserves at the district level initially and latter stages from the near by districts. Hay should be enriched with 2% Urea molasses solution or 1% common salt solution and fed to LS</p> <p>Herd should be split and supplementation should be given only to the highly productive and breeding animals</p> <p>Provision of emergency grazing/feeding (Cow-calf camps or other special arrangements to protect high productive &amp; breeding stock)</p> <p>Motivate the farmers to mix the dry fodder with available kitchen waste while feeding</p> <p>Arrangements should be made for</p>	<p>Concentrates supplementation should be provided to all the animals.</p> <p>The farmers may be advised to practice “flushing the stock” to recoup</p> <p>Short duration fodder crops of should be sown in unsown and crop failed areas where no further routine crop sowing is not possible</p> <p>Supply of quality seeds of fodder varieties and motivating the farmers to cultivate at least 10% of their land holding for fodder production</p>

	<p>good quality chaff cutters.</p> <p>Harvesting and collection of perennial vegetation particularly grasses which grow during monsoon</p> <p>Proper drying, bailing and densification of harvested grass from previous season</p> <p>Creation of permanent fodder, feed and fodder seed banks in all drought prone villages</p>	<p>mobilization of small ruminants across the villages where no drought exits with subsidized road/rail transportation and temporary shelter provision for the shepherds</p> <p>Unproductive livestock should to be culled during severe drought</p> <p>Create transportation and marketing facilities for the culled and unproductive animals</p> <p>Supply silage and or hay on subsidized rates to the farmers having high productive stock</p> <p>Subsidized loans should be provided to the livestock keepers</p>	
<b>Cyclone</b>	<p>Harvest all the possible wetted grain (rice/bajra/maize/greengram/blackgram/groundnut etc) and sugar cane tops and use as animal feed.</p> <p>Motivate the farmers to store a minimum quantity of hay (25-50 kg) and concentrates (10-25 kg) per animal in farmer's/LS keepers house/shed for feeding the animals during cyclone.</p> <p>Stock of anti-diarrheal drugs and electrolytes should be made available for emergency transport</p> <p>Don't allow the animals for grazing in case of early forewarning (EFW) of cyclone</p> <p>Incase of EFW of severe cyclone, shift the animals to safer places.</p>	<p>Treatment of the sick, injured and affected animals through arrangement of mobile emergency veterinary hospitals/rescue animal health workers.</p> <p>Diarrhea out break may happen. Health camps should be organized</p> <p>In severe cases un-tether <b>or</b> let loose the animals</p> <p>Arrange transportation of highly productive animals to safer place</p> <p>Spraying of fly repellants in animal sheds</p>	<p>Repair of animal shed</p> <p>Deworm the animals through mass camps</p> <p>Vaccinate against possible disease out breaks like HS, BQ, FMD and PPR</p> <p>Proper dispose of the dead animals / carcasses by burning / deep burying (4-8 feet) with lime powder (1kg for small ruminants and 5kg for large ruminants) in pit</p> <p>Bleach / chlorinate (0.1%) drinking water or water resources</p> <p>Collect drowned crop material, dry it and store for future use</p>

			<p>Sowing of short duration fodder crops in unsown and water logged areas when crops are damaged and no chance to replant</p> <p>Application of urea (20-25kg/ha) in the inundated areas and CPR's to enhance the bio mass production.</p>
<b>Floods</b>	<p>In case of early forewarning (EFW), harvest all the crops (rice/maize/greengram/blackgram) that can be useful as fodder in future (store properly) and also sugar cane tops</p> <p>Don't allow the animals for grazing if floods are forewarned</p> <p>Arrangement for transportation of animals from low lying area to safer places and also for rescue animal health workers to get involve in rescue operations</p>	<p>Transportation of animals to elevated areas</p> <p>Stall feeding of animals with stored hay and concentrates</p> <p>Proper hygiene and sanitation of the animal shed</p> <p>In severe floods, un-tether or let loose the animals</p> <p>Emergency outlet establishment for required medicines or feed in each village</p> <p>Spraying of fly repellants in animal sheds</p>	<p>Repair of animal shed</p> <p>Bring back the animals to the shed</p> <p>Cleaning and disinfection of the shed</p> <p>Bleach (0.1%) drinking water / water sources</p> <p>Deworming with broad spectrum dewormers</p> <p>Vaccination against possible disease out breaks like HS, BQ, FMD and PPR</p> <p>Proper disposable of the dead animals / carcasses by burning / deep burying (4-8 feet) with lime powder (1kg for small ruminants and 5kg for large ruminants) in pit</p> <p>Drying the harvested</p>



			crop material and proper storage for use as fodder.
--	--	--	---

**Vaccination programme for cattle and buffalo:**

<b>Disease</b>	<b>Age and season at vaccination</b>
Anthrax	In endemic areas only, Feb to May
Haemorrhagic septicaemia (HS)	May to June
Black quarter (BQ)	May to June
Foot and mouth disease (FMD)	July/August and November/December

**Vaccination schedule in small ruminants (Sheep & Goat)**

<b>Disease</b>	<b>Season</b>
Foot and mouth disease (FMD)	Preferably in winter / autumn
Peste des Petits Ruminants (PPR)	Preferably in January
Black quarter (BQ)	May / June
Enterotoxaemia (ET)	May
Haemorrhagic septicaemia (HS)	March / June
Sheep pox (SP)	November

### 2.5.2 Poultry

	Suggested contingency measures		
	Before the event <sup>a</sup>	During the event	After the event
<b>Drought</b>			
Shortage of feed ingredients	Storing of house hold grain like maize, broken rice, etc, in to use as feed in case of severe drought	Supplementation only for productive birds with house hold grain Supplementation of shell grit (calcium) for laying birds Culling of weak birds	Supplementation to all survived birds
Drinking water		Use water sanitizer or offer fresh and cool drinking water	
Health and disease management	Culling of sick birds. Deworming and vaccination against RD and fowl pox	Mixing of Vit. A,D,E, K and B-complex including vit C in drinking water (5ml in one litre water)	Hygienic and sanitation of poultry house Disposal of dead birds by burning / burying with lime powder in pit
<b>Floods</b>			
Shortage of feed ingredients	In case of early forewarning of floods, shift the birds to safer place Storing of house hold grain like maize, broken rice, etc,	Use stored feed as supplement Don't allow for scavenging Culling of weak birds	Routine practices are followed Deworming and vaccination against RD
Drinking water		Use water sanitizer or offer fresh drinking water	
Health and disease management	In case of EFW, add antibiotic powder (Terramycin/Ampicilline/ Ampiclox etc., 10g in one litre) in drinking water to prevent any disease outbreak	Prevent water logging surrounding the sheds through proper drainage facility Assure supply of electricity by generator or solar energy or biogas Sprinkle lime powder to prevent ammonia accumulation due to dampness	Sanitation of poultry house Treatment of affected birds Disposal of dead birds by burning / burying with lime powder in pit Disposal of poultry manure to prevent protozoal problem Supplementation of coccidiostats in feed Vaccination against RD
<b>Cyclone</b>			

Shortage of feed ingredients	In case of EFW, shift the birds to safer place Storing of house hold grain like maize, broken rice, bajra etc, Culling of weak birds	Use stored feed as supplement Don't allow for scavenging Protect from thunder storms	Routine practices are followed
Drinking water		Use water sanitizers or offer cool drinking water	
Health and disease management	In case of EFW, add antibiotic powder in drinking water to prevent any disease outbreak	Sanitation of poultry house Treatment of affected birds Prevent water logging surrounding the sheds Assure supply of electricity Sprinkle lime powder (5-10g per square feet) to prevent ammonia accumulation due to dampness	Disposal of dead birds by burning /deep burying with lime powder in pit Disposal of poultry manure to prevent protozoal problem Supplementation of coccidiostats in feed Vaccination against Ranikhet Disease (0.5ml S/c)
<b>Heat wave and cold wave</b> NA			

<sup>a</sup> based on forewarning wherever available

<b>Andhra Pradesh Contingency plans for FISHERIES/AQUACULTURE</b>			
	<b>Suggested Contingency Measures</b>		
<b>1) Drought</b>	<b>Before the event</b>	<b>During the event</b>	<b>After the event</b>
<b>A. Capture</b>			
Marine	No intervention	No intervention	No intervention
Inland			

(i) Shallow water depth due to insufficient rains / inflow	Stocking of advanced fingerlings in half or even less than the normal stocking density or stocking of common carp seed	Immediate harvesting or decreasing the density commensurate with the water quantity.	De weeding and deepening of tank to ensure retention of water for a longer period and provision of employment under MGNREGP
(ii) Changes in water quality	Regular monitoring of water quality parameters and application of geolites, soil probiotics, etc to maintain water quality	Immediate harvesting or changing the water quality by application of sanitisers.	Removal of top layer, deep ploughing of tank and application of lime
(iii) Any other			
<b>B. Aquaculture</b>			
(i) Shallow water in ponds due to insufficient rains / inflow	Crop holiday or going for stocking of yearlings by reducing the density according to availability of water	Harvesting of fish and leaving the pond fallow till next season	Removal of top layer, deep ploughing of tank and application of lime
(ii) Impact of salt load build up in ponds /changes water quality	Stocking of salinity tolerant fish/ shrimp, application of geolites and other buffers	Frequent change of water with fresh water	Frequent draining of the pond with fresh water, removal of top layers
(iii) Any other			
<b>2) Floods</b>			
<b>A. Capture</b>			
Marine	No intervention	No intervention	No intervention
Inland			
(i) Average compensation paid due to loss of human life	Shifting the people from low lying areas to relief camps	Deployment of specially trained persons for rescue operations by providing life bouys, jackets, ropes, boats, etc	Payment sufficient ex-gratia to the families

(ii) No. of boats / nets damaged	Shifting and relocating boats and nets to safer places when warnings are issued, to avoid fishing, etc	Shifting and relocating boats and nets to safer places	Assessment of damages to boats and nets and provision of boats and nets for restoration of livelihoods
(iii) No. of houses damages	Avoidance of construction of houses in flood prone areas, construction of pucca houses at elevated places,	Shifting of people by relief boats to the relief camps	Assessment of damages to houses and provision of compensation in case of partial damage and sanction house under existing schemes
(iv) Loss of stock	Avoidance of surface species like catla, silver carp since they are vulnerable in tanks prone to floods, erection of nets across the spill way or just beyond it	Erection of nets at spill ways	Taking up compensatory stocking
(v) Changes in water quality		When dissolved oxygen levels go down, aerators, recirculation of water, etc are to be attempted to maintain DO levels, going for partial harvest, etc	
(vi) Health and disease	Sometimes there may be heavy accumulation of nutrients and organic matter.	There may be break out of Hemorrhagic septicemia. Addition of antibiotics like Chloro Tetra Cycline or Oxy Tetra Cycline to the feed to control the disease	Removal of weeds, top layer of soil, deep ploughing of tank and application of lime, exposing to sun light
<b>B. Aquaculture</b>			
(i) Inundation with flood water	Raising and rivetting the bunds, construction of spill way to release excess water, erection of nets to avoid escape of fish	Continuous pumping of excess water, erection of nets low lying areas	Strengthening of bunds, excavating channels along the sides of the ponds for free escape of water

(ii) Water contamination and changes in water quality		When dissolved oxygen levels go down, aerators, recirculation of water, etc are to be attempted to maintain DO levels, going for partial harvest, etc	
(iii) Health and disease	Sometimes there may be heavy accumulation of nutrients and organic matter.	There may be break out of Hemorrhagic septicemia. Addition of antibiotics like Chloro Tetra Cycline or Oxy Tetra Cycline to the feed to control the disease	Removal of weeds, top layer of soil, deep ploughing of tank and application of lime, exposing to sun light
(iv) Loss of stock and inputs (feed, chemicals, etc)	Advance erection of nets, strengthening of bunds where they are prone to breaches, harvesting or reducing the density	Suspension of feeding, application of organic manures	Compensatory stocking, assessment of values and payment of subsidy on inputs
(v) Infrastructure damage (pumps, aerators, huts, etc.)	Insuring pond, accessories, etc., Shifting of aerators, pumps soon after warnings are issued	Relocating pumps, aerators to elevated places	Assessment of damages and provision of them on subsidy
(vi) Any other			
<b>3) Cyclone /Tsunami</b>			
<b>A. Capture</b>			
Marine			
(i) Average compensation paid due to loss of fishermen lives	Avoidance of fishing, preventing fishermen from venturing into sea, carrying of safety equipment and VHF sets, shifting fishermen from vulnerable areas to relief camps, etc	To ensure the return of fishing boats on long voyages, provision of information on such boats to coast Guard	Payment sufficient ex-gratia to the families

(ii) Average no. of boats / nets damaged	Avoidance of fishing when warnings are issued, shifting of boats and nets to safe places	Shifting and relocating boats and nets to safer places	Assessment of damages to boats and nets and provision of boats and nets for restoration of livelihoods
(iii) Average no. of houses damages	Avoidance of houses in Coastal Regulation Zone, designing of houses to withstand impact of turbulent wind and water	Shifting of people by relief boats to the relief camps	Assessment of damages to houses and provision of compensation in case of partial damage and sanction house under existing schemes
Inland	Erection of protective nets across the surplus weir to prevent fish loss due to overflows	Continuous monitoring to prevent or minimise escape of fish along with surplus water	Compensatory stocking of seed
<b>B. Aquaculture</b>			
(i) Overflow / flooding in ponds	The design of the pond must be in such a manner as to bail out surplus water and to prevent loss of standing crop	Continuous monitoring to prevent or minimise escape of fish along with surplus water	Compensatory stocking of seed
(ii) Changes in water quality (fresh water / brackish water ratio)	Recirculation water to replenish and ensure sufficient dissolved oxygen levels in the pond. Maintenance of salinity levels by pumping in water from creeks.	Continuation of the same process.	Restoration of physical and chemical parameters
(iii) Health and disease	Removal of stress causing factors to maintain the health of the animal	Removal of stress causing factors to maintain the health of the animal	Restoration of physical and chemical parameters
(iv) Loss of stock and inputs (feed, chemicals, etc)	Preventive nets must be erected to minimise loss of stock	Continuation of the same process.	Compensatory stocking of seed

(v) Infrastructure damage (pumps, aerators, huts, etc.)	Pumps, aerators, etc must be protected by moving them to safe locations	To avoid use of aerators, pumps and other appliances	Overhauling of the equipment to prevent from being damaged
(vi) Any other			
<b>4) Heat and Cold wave conditions</b>			
<b>A. Capture</b>			
Marine	Avoidance of fishing	Avoidance of fishing	No intervention
Inland	Monitoring dissolved oxygen levels	Monitoring dissolved oxygen levels	No intervention
<b>B. Aquaculture</b>			
(i) Changes in water quality (fresh water /brackish water ratio)	Reduction of biomass by partial harvest in the event of heat as the DO levels will be very low.	Avoidance of fishing	Compensatory stocking of seed and restoration of all physical and chemical parameters
(ii) Health and disease	Removal of stress causing factors to maintain the health of the animal	Removal of stress causing factors to maintain the health of the animal	Compensatory stocking of seed and restoration of all physical and chemical parameters
(iii) Any other			



# Contingency Plans for Rabi Crops for Krishna Zone

## Prakasam District

### 1. Rainfall Information

	Oct – Dec	Jan – Mar
(a) Normal rainfall during <i>Rabi</i> season	: 393	-
(b) Number of rainy days	: 18	-

### 2. Rabi crops cultivated

#### 2aArea Production statistics

S. No	Cropping System	Crop name	Area ‘000 ha	Production ‘000 t	Productivity Kg/ha
1		Paddy	37490	158633	4231
2		Jowar	20714	3704	179
3		Bajra	31	25	803
4		Maize	4267	17440	4087
5		Ragi	379	475	1254
6		Other Minor millets	2204	1730	785
7		Total Coarsegrains	27595	23374	-
8		Bengal gram	109283	74640	683
9		Redgram	314	60	191
10		Greengram	647	211	326
11		Blackgram	14890	3991	268
12		Horsegram	1398	629	450
13		Cowgram	10389	5195	500
14		Other Pulses	0	0	0
15		Total Pulses	136921	84725	-
16		Total Foodgrains	202006	266733	-

17		Groundnut	855	1261	1475
18		Sesamum	2702	211	78
19		Sunflower	618	158	255
20		Safflower	0	0	0
21		Castor	1132	274	242
22		Soyabean	0	0	0
23		Niger	0	0	0
24		Rapeseed and Mustard	9	5	500
25		<b>Other Oilseeds</b>	<b>0</b>	<b>0</b>	<b>0</b>
26		<b>Total Oilseeds</b>	<b>5316</b>	<b>1908</b>	<b>-</b>
27		Cotton (kapas)	255	237	930
28		Mesta	0	0	0
29		Chillies	5127	8885	1733
30		Sugarcane	0	0	0
31		Onion	11	83	7500
32		Turmeric	0	0	0
33		Tobacco	45509	53276	1171
34		Coriander	117	76	650
		Total cropped area	258341	331197	-

2b. Source wise (Water) cultivated area

Sources of Irrigation	Number	Area ('000 ha)	Percentage of total irrigated area
Canals		66.7	34.3
Tanks	957	27.8	14.3
Open wells	22783		
Bore wells	41163	84.4	43.4
Lift irrigation	1407		
Micro-irrigation	--		
Other sources	--	15.7	8.1
Total Irrigated Area		194.6	100.0

3. Sowing window information

S. No.	Soil type	Cropping system	Crop name	Optimum sowing window (Please mention along with week i.e., 2 <sup>nd</sup> week of Oct-4 <sup>th</sup> week of Nov/etc.)
1	Black soils	Fallow/Korra-Bengalgram	Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov
2		Fallow-Blackgram	Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov
3		Fallow-Sesamum	Sesamum	2 <sup>nd</sup> Week of Dec to 1 <sup>st</sup> week of Jan
4		Fallow-Coriander	Coriander	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov
5		Fallow-Redgram	Rabi Redgram	1st week of Oct to 2 <sup>nd</sup> week of Oct
6		Fallow-Tobacco	Natu Tobacco	

7			Cigarette tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov
8		Fallow-Maize Greengram-Maize	Maize	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov
9		Fallow-Sunflower	Sunflower	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov
11	Red soils	Fallow-Blackgram	Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov
12		Fallow-Greengram	Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov
13		Fallow-Cowpea	Cowpea	2 <sup>nd</sup> week of Oct to 1 <sup>st</sup> week of Nov
14		Fallow-Castor	Castor	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov
15		Fallow-Jowar	Jowar	1 <sup>st</sup> week of Oct to 1 <sup>st</sup> week of Nov
16		Fallow - Tobacco	Natu Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov
17		Fallow - Tobacco	FCV tobacco	

#### 4. Contingency measures Field crops

**For crops grown with residual moisture i.e., under rainfed condition**

(a) Excess residual moisture

S. No.	Soil type	Cropping system	Crop name	Sowing Window	Variety	Management practices
1	Rainfed black soil	Fallow-Bengalgram	Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-	Normal practices and need based plant protection

					399	measures
2		Fallow-Maize  Blackgram-Maize	Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January	DHM-115, 117, KH-510  Private hybrids	Normal practices and need based plant protection measures
3		Fallow-Jowar	Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Normal practices and need based plant protection measures
4		Fallow-Blackgram	Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	Normal practices and need based plant protection measures
5		Fallow-Greengram	Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM-2-14,	Normal practices and need based plant protection measures
6		Rice-Sesamum  Fallow-Sesamum	Sesamum	3 <sup>rd</sup> week of December to 1 <sup>st</sup> week of January	YLM-11, 17, 66,	Normal practices and need based plant protection measures
7		Fallow-Tobacco	Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	Siri, VT-1135, G-11 & CH-3	Normal practices and need based plant protection measures
8		Fallow -Variga	Variga	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov	Nagarjuna, Sagar, Co-4	Normal practices and need based plant protection measures
9		Fallow-Rabi redgram	Rabi redgram	1 <sup>st</sup> week of Oct to 2 <sup>nd</sup> week of Oct	LRG-52, LRG-41,BSMR736	Reduced spacing of redgram from 180 cm to 90 cm

10	Red soils	Fallow-Greengram	Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM-2-14,	Normal practices and need based plant protection measures
11		Fallow-Cowpea	Cowpea	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TPTC-29, C 152, CO-8, APFC-10/1	Normal practices and need based plant protection measures
12		Fallow-Castor	Castor	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	PCH-111, PCH-222, Kranthi, Haritha	Normal practices and need based plant protection measures
13		Fallow-Tobacco	Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	Siri, VT-1135, G-11 & CH-3	Normal practices and need based plant protection measures
14			Variga	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov	Nagarjuna, Sagar, Co-4	Normal practices and need based plant protection measures
15	Sandy loams	Groundnu-Groundnut	Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K6, K9, Kadiri Harithandra, Dharani	Foliar spray of urea 2 % or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> and FeSO <sub>4</sub> spray; Spraying of copper oxy chloride or mancozeb etc. to manage fungal diseases.

(b) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
--------	-----------	-----------------	-----------	-------------	---------	----------------------

1	Black soil	Fallow - Bengalgram	Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-399	1. Need based application of pesticides to manage insect pests and diseases, 2. Inter cultivation practices 3. Life saving irrigation with micro irrigation systems at critical growth stages 4. Conservation furrow methods 5. Foliar spray of 2% urea Or 1% KNO <sub>3</sub>
2		Fallow-Jowar	Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	
3		Fallow-Blackgram	Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	
4		Fallow-Greengram	Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM-2-14,	
5		Rice-Sesamum  Fallow-Sesamum	Sesamum	3 <sup>rd</sup> week of December to 1 <sup>st</sup> week of January	YLM-11, 17, 66,	
6		Fallow-Tobacco	Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	Siri, VT-1135, G-11 & CH-3	
7		Fallow - Variga	Variga	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov	Nagarjuna, Sagar, Co-4	
8	Red soil	Fallow-Rabi greengram	Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM-2-14,	
9		Fallow-Castor	Cowpea	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TPTC-29, C 152, CO-8, APFC-10/1	

10		Fallow-Tobacco	Castor	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	PCH-111, PCH-222, Kranthi, Haritha	
11		Fallow-Tobacco	Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	Siri, VT-1135, G-11 & CH-3	
12		Fallow - Variga	Variga	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov	Nagarjuna, Sagar, Co-4	

(c) Severe limitation in moisture. Deficit of rainfall during September/October months by more than 40%.

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Black soil		Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-399	1. Need based application of pesticides to manage insect pests and diseases, 2. Inter cultivation practices 3. Life saving irrigation with micro irrigation systems at critical growth stages 4. Conservation furrow methods 5. Foliar spray of 2% urea Or 1% KNO <sub>3</sub>
2			Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	
3			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	
4			Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM-2-14,	
5			Sesamum	3 <sup>rd</sup> week of December to 1 <sup>st</sup>	YLM-11, 17, 66,	



				week of January		
6			Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	Siri, VT-1135, G-11 & CH-3	
7			Variga	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov	Nagarjuna, Sagar, Co-4	
8	Red soil		Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM-2-14,	
9			Cowpea	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TPTC-29, C 152, CO-8, APFC-10/1	
10			Castor	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	PCH-111, PCH-222, Kranthi, Haritha	
11			Tobacco	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	Siri, VT-1135, G-11 & CH-3	
12			Variga	2 <sup>nd</sup> week of Oct to 3 <sup>rd</sup> week of Nov	Nagarjuna, Sagar, Co-4	

### For crops grown with groundwater

(a) Above normal rainfall in *Kharif* coupled with good distribution

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Black soils/Red	Rice-Rice	Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup>	NLR-34449	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage

	soils			Week of Nov	NLR-3042 NLR-4001 JGL-384 MTU-1156 MTU-1153 MTU-1121	insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
2	Red soils	Rice-Sesamum  Fallow-Sesamum	Sesamum	3 <sup>rd</sup> week of December to 1 <sup>st</sup> week of January	YLM-17, 66	
3	Sandy loams	G.Nut-G.nut	Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition

(b) Normal rainfall

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Black soil		Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, KAK-2, Vihar, NBeG-399	

2			Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4 NJ-2647, CSH-9, 14, 16,	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
3			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	
4			Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January	DHM-115, 117, KH-510  Private hybrids	
5			Sesamum	3 <sup>rd</sup> week of December to 1 <sup>st</sup> week of January	YLM-11, 17, 66,	
6			Rabi redgram	2 <sup>nd</sup> week of Oct to 2 <sup>nd</sup> week of Nov	LRG-52, LRG-41	

(c) Deficient rainfall in *Kharif* season (25-50% deficient)

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
1	Black soil		Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, KAK-2, Vihar,	Foliar spray  Life saving irrigation through micro irrigation system

					NBeG-399	
2			Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4 NJ-2647, CSH-9, 14, 16,	
3			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU- 31, LBG- 752	
5			Sesamum	3 <sup>rd</sup> week of December to 1 <sup>st</sup> week of January	YLM-11, 17, 66,	

(d) Scanty rainfall in *Kharif* season

S. No.	Soil type	Cropping system	Crop name	Sowing time	Variety	Management practices
	Rain black soils/ red soils	Fallow- Bengalgram/ Jowar/ blackgram/ Cowpea/ Sesamum	Horsegram/	4 <sup>th</sup> week of Sept	CRIDA 18-R  CRIDA-22B	1.Foliar spray with 2% Urea  2.Life saving irrigation through micro irrigation system
			Greengram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TM-96-2, WGG-42, IPM- 2-14,	
			Fodder jowar			

• **Management practices for unseasonal rains**

Condition - Continuous high rainfall in a short span leading to water logging and Heavy rainfall with high speed winds in a short span				
Crop	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Rice	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>• Take up gap filling either with available nursery or by splitting the tillers from the surviving hills</li> <li>• Take up suitable plant protection Measures in anticipation of pest &amp; disease out breaks</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>• Take up suitable plant protection Measures in anticipation of pest &amp; disease out breaks</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Take up suitable plant protection measures in anticipation of pest &amp; disease out breaks</li> </ul>	<ul style="list-style-type: none"> <li>• Drain out water and spread sheaves loosely in field or field bunds where there is no water stagnation</li> <li>• spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds</li> <li>• Thresh after drying the sheaves properly Ensure proper grain moisture before storing</li> </ul>
Blackgram/ Greengram/ Mungbean/ Cowpea/ Soybean	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 4-5 kg N/ha after draining excess water</li> <li>• Spray KNO<sub>3</sub> 1% or water soluble fertilizers like 19-19-19,20-20-20, 21-21-21 at 1% to support nutrition</li> <li>• Spray fungicides like Copper oxy chloride 0.3 % or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</li> <li>• Take up timely control measures against the outbreak of pests like Spodoptera etc.</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 4-5 kg N /ha after draining excess water</li> <li>• Spray KNO<sub>3</sub> 1% or water soluble fertilizers like 19-19-19,20-20-20, 21-21-21 at 1% to support nutrition</li> <li>• Spray fungicides like Copper oxy chloride 0.3% or Carbendazim 0.1 % or Mancozeb 0.25% two to three times by rotating the chemicals</li> <li>• Take up timely control measures against the outbreak of pests like Spodoptera etc.</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Allow the crop to dry completely before harvesting</li> </ul>	<ul style="list-style-type: none"> <li>• Spread the bundles drenched in rain on field bunds or drying floors to quicken the drying</li> <li>• Thresh the bundles after they are dried properly Dry the grain to proper</li> <li>• moisture per cent before</li> <li>• bagging and storing to prevent deterioration in quality during storage</li> </ul>

Maize	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>• Take up inter cultivation and at optimum soil moisture condition to loosen and aerate the soil and to control weeds</li> <li>• Earthen up the crop for anchorage Spray KNO<sub>3</sub> 1% or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</li> <li>• Take up timely control measures for Pink stem borer, sheath blight and Turcicum leaf blight</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Apply 20 kg N + 10 kg K /ha after draining excess water</li> <li>• Spray KNO<sub>3</sub> 1% or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</li> <li>• Take up timely control measures for sheath blight and post flowering stalk rots</li> </ul>	<ul style="list-style-type: none"> <li>• Drain the excess water as early as possible</li> <li>• Allow the crop to dry completely before harvesting</li> </ul>	<ul style="list-style-type: none"> <li>• Harvest the cobs after the they are dried up properly. Dry the grain to optimum moisture condition before storing</li> </ul>
-------	--	---	---	---

**For crops grown with Canal Irrigation: The scenario would be based on the storage available in the reservoirs.**

a. Limited release of water

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soil & Red soils	Rice – Rice	Maize	3 <sup>rd</sup> week of November to 1 <sup>st</sup> week of January	DHM-115, 117, KH-510 Private hybrids	Foliar spray of urea 2% or KNO <sub>3</sub> 1%, ZnSO <sub>4</sub> and FeSO <sub>4</sub> spray; Spraying of copper oxy chloride or mancozeb etc. to manage fungal diseases. Intercultivation to reduce soil moisture
			Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	NLR-34449 NLR-3042 NLR-4001 JGL-384 MTU-1156 MTU-1153	Direct sowing with drum seeder

					MTU-1121	
2			Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
3			Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-399	
4			Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
5			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	

b. Delayed release of water

**Head reach:**

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soils/Red soils	Rice-Rice	Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	NLR-34449 NLR-3042	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages

					NLR-4001 JGL-384 MTU-1156 MTU-1153 MTU-1121	or based on soil moisture condition
--	--	--	--	--	---	-------------------------------------

**Middle reach:**

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soils/Red soils	Rice-Rice	Rice	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	NLR-34449 NLR-3042 NLR-4001 JGL-384 MTU-1156 MTU-1153 MTU-1121	Direct sowing with drum seeder



**Tail end:**

S. No.	Soil type	Cropping system	Crop name	Sowing window	Variety	Management practices
1	Black soil & Red soils	Rice – Rice	Jowar	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	NTJ-2,4, NJ-2647, CSH-9, 14, 16,	Seed treatment, Pre and post emergence application of weedicides, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
2			Bengalgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> Week of Nov	JG-11, NBeG-49, NBeG-47, KAK-2, Vihar, NBeG-399	Seed treatment, Pre and post emergence application of weedicides, gypsum application, need based application of pesticides to manage insect pests and diseases, irrigation at critical growth stages or based on soil moisture condition
3			Groundnut	2 <sup>nd</sup> week of November to 2 <sup>nd</sup> week of December	TAG 24, K-6, K-9, Kadiri Harithandra, Dharani	
			Blackgram	2 <sup>nd</sup> week of Oct to 4 <sup>th</sup> week of Nov	TBG-104, GBG-1, PU-31, LBG-752	

**5. Contingency measures for Horticulture Crops (Existing / New plantations)**

S. No.	Crop Name	Specific management practices to be taken up following excess/deficient/scanty rainfall	Time of intervention	Remarks

**6. Contingency measures for Horticulture Crops (vegetables)**

S. No.	Crop Name	Specific management practices to be taken up following excess/deficient/scanty rainfall	Time of intervention	Remarks

**7. Temperature related stresses for field and horticulture crops:**

Excess temperatures/Less than normal temperatures

S.No	Crop name	Stage of crop growth	Threshold temperature*	Suggested management practices


\* Temperature increase or decrease over normal and for number of days . For example, increase of 3 degrees over normal for a period of 5 days

**8. Management practices for livestock** (to cover shelter management during cold or heat waves, production/regulation of fodder in rabi season in deficient monsoon years/ excess monsoon rainfall years etc),

**For Fodder crops grown with residual moisture i.e., under rainfed condition**

(a) Excess (rainfall during September/October months) residual moisture

S. No.	Soil type	Cropping system	Fodder name	Variety	Management practices

(b) Normal rainfall (rainfall during September/October months) residual moisture

S. No.	Soil type	Cropping system	Crop name	Variety	Management practices

(b) Less than optimum moisture i.e., 25% less than normal, which can happen due to insufficient rainfall during September/October months. Deficit of 20-40% rainfall

S. No.	Soil type	Cropping system	Fodder name	Variety	Management practices

(c) Severe limitation in moisture. Deficit of rainfall during September/October months by more than 40%.

S. No.	Soil type	Cropping system	Fodder name	Variety	Management practices

**For fodder crops (mostly perennial fodder varieties as sole fodder crop) grown with groundwater**

S. No.	Soil type	Fodder name	Variety	Management practices

**Livestock management during severe cold waves/heat waves**

Nutritional management	Shelter management	Health management	Miscellaneous, if any