

State: Arunachal Pradesh
Agriculture Contingency Plan: East Siang District

| 1.0 District Agriculture profile* | | | | |
|--|--|---|---|------------------|
| 1.1 | Agro-Climatic/Ecological Zone | 16.3 Arunachal Pradesh (Subdued Eastern Himalayas), warm to hot, perhumid eco-subregion (C1A10) | | |
| | Agro Ecological Sub Region (ICAR) | Humid Bengal – Assam Basin | | |
| | Agro-Climatic Zone (Planning Commission) | Eastern Himalayan Region | | |
| | Agro Climatic Zone (NARP) | North East Hill Region | | |
| | List all the districts falling under the NARP Zone* (*>50% area falling in the zone) | East Siang | | |
| | Geographic coordinates of district headquarters | Latitude | Longitude | Altitude |
| | | 27 ^o 43' - 29 ^o 20' N | 94 ^o 42' - 95 ^o 35' E | 151 – 4000 m MSL |
| | Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS | ICAR Research Complex for NEH Region, AP Centre, Basar, West Siang district, Arunachal Pradesh ICAR Research Complex for NEH Region, Umroi Road, Barapani, Meghalaya | | |
| | Mention the KVK located in the district with full address | KVK East Siang, CHF, CAU, Pasighat, Arunachal Pradesh-791102 | | |
| Name and address of the nearest Agromet Field Unit (AMFU, IMD) for agro-advisories in the Zone | ICAR Research Complex for NEH Region, AP Centre, Basar, West Siang district, Arunachal Pradesh | | | |

***Indicate source of data while furnishing information at different places in the district profile**

| 1.2 | Rainfall | Normal RF (mm) | Normal Rainy days (number) | Normal Onset (specify week and month) | Normal Cessation (specify week and month) |
|-----|-----------------------------------|----------------|----------------------------|---------------------------------------|---|
| | Monsoon/SW monsoon (June-Sep) | 2548.4 | 67 | 1 st week June | Last week of September |
| | Post monsoon/NE monsoon (Oct-Dec) | 232.4 | 11 | 1 st week October | Last week of October |
| | Winter (Jan- February) | 173.6 | 19 | 3 rd week February | Last week of March |
| | Pre monsoon/Summer (April-May) | 779.2 | 31 | 1 st week April | Last Week of May |
| | Annual | 3733.6 | 128 | - | - |

| 1.3 | Land use pattern of the district | Geographical area | Cultivable area | Forest area | Land under non-agricultural use | Permanent pastures | Cultivable wasteland | Land under Misc. tree crops and groves | Barren and uncultivable land | Current fallows | Other fallows |
|-----|----------------------------------|-------------------|-----------------|-------------|---------------------------------|--------------------|----------------------|--|------------------------------|-----------------|---------------|
| | Area ('000 ha) | 400.50 | 40.92 | 227.85 | 2.53 | 1.09 | 6.66 | 3.17 | 3.33 | 3.01 | 3.4 |

2011 Stats Directorate of Economics and Statistics, Ministry of Agriculture, Govt. of India

| 1.4 | Major Soils (common names like red sandy loam deep soils (etc.,))* | Area ('000 ha)** | Percent (%) of total geographical area |
|-----|--|------------------|--|
| | 1. Red Soil | 47.40 | 11.83 |
| | 2. Sandy Soil | 107.90 | 26.94 |
| | 3. Sandy Loam | 132.60 | 33.10 |
| | 4. Loamy Sand | 82.40 | 20.57 |
| | 5. Others | 30.20 | 7.54 |

| |
|--|
| Soil pH 4.3 – 6.8, Organic Carbon: 0.11% - 3.90% Low to Medium to High |
|--|

(Source: Statistical Abstract, Arunachal Pradesh, 2007)

| 1.5 | Agricultural land use | Area ('000 ha) | Cropping intensity % |
|------------|------------------------------|-----------------------|-----------------------------|
| | Net sown area | 24.69 | 126.6% |
| | Area sown more than once | 6.56 | |
| | Gross cropped area | 31.25 | |

(Source: 2011-12 Stats Directorate of Economics and Statistics, Ministry of Agriculture, Govt. of India)

| 1.6 | Irrigation | Area ('000 ha) | | |
|------------|--|-------------------------------|-----------------------|--|
| | Net irrigated area | 11.67 | | |
| | Gross irrigated area | 11.67 | | |
| | Rainfed area | 13.87 | | |
| | Sources of Irrigation | Number | Area ('000 ha) | Percentage of total irrigated area |
| | Stream flow | 2 | 11.63 | 99% |
| | Tanks | - | - | - |
| | Open wells | - | - | - |
| | Bore wells | - | - | - |
| | Lift irrigation schemes | - | - | - |
| | Micro-irrigation | - | - | - |
| | Other sources (please specify) | - | - | 1% |
| | Total Irrigated Area | - | 11.67 | 100% |
| | Pump sets | 353 | - | - |
| | No. of Tractors | 86 | - | - |
| | Groundwater availability and use* (Data source: State/Central Ground water Department /Board) | No. of blocks/ Tehsils | (%) area | Quality of water (specify the problem such as high levels of arsenic, fluoride, saline etc) |
| | Over exploited | - | - | - |
| | Critical | - | - | - |
| | Semi- critical | - | - | - |
| | Safe | 6 | 100 | - |
| | Wastewater availability and use | - | - | - |
| | Ground water quality | - | | |

*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%

| 1.6. a. | Fertilizer and Pesticides use | Type | Total quantity (MT) |
|---------|-------------------------------|---|----------------------------------|
| 1 | Fertilizers* | Nitrogen (N) Phosphate (P ₂ O ₅) Potassic (K ₂ O) Total NPK Other straight fertilizers (specify) Other complex fertilizers (specify) | 43.70 18.50 10.50 72.70 |
| 2 | Chemical Pesticides* | Insecticides Fungicides Weedicides Others (specify) | |

(Source: Statistical Abstract, Arunachal Pradesh, 2007)

For year 2007-08:

Fertilizer in (Ton) N: 26.67; P: 25.81; K: 44.82

Pesticides (Lit) Monocrotophos/Roger: 336.5

(Source: Respective Department of East Siang district, Arunachal Pradesh, 2008)

1.7 Area under major field crops & horticulture (2006-07)

| 1.7 | S. No. | Major field crops cultivated | Area ('000 ha) | | | | | | | Grand total |
|-----|---------|------------------------------|----------------|---------|-------|-------------|---------|-------|--------|-------------|
| | | | <i>Kharif</i> | | | <i>Rabi</i> | | | Summer | |
| | | | Irrigated | Rainfed | Total | Irrigated | Rainfed | Total | | |
| 1 | Paddy | - | 13.137 | 13.137 | - | - | - | - | 16.770 | |
| 2 | Maize | - | 2.883 | 2.883 | - | - | - | - | 4.050 | |
| 3 | Millet | - | 2.220 | 2.220 | - | - | - | - | 2.461 | |
| 4 | Oilseed | - | - | - | - | 1.735 | 1.735 | - | 5.075 | |
| 5 | Potato | - | - | - | 0.294 | - | 0.294 | - | 0.875 | |
| 6 | Pulses | - | 0.937 | 0.937 | - | - | - | - | 1.028 | |

| | | | | | | | | | | |
|--|---|--------|---|-------|-------|---|---|---|---|------|
| | 7 | Ginger | - | 0.177 | 0.177 | - | - | - | - | 0.98 |
|--|---|--------|---|-------|-------|---|---|---|---|------|

NEDFI Databank website, 2012-13 Statistical Abstract of Arunachal Pradesh.

| | S. No. | Horticulture crops Fruits | Area ('000 ha) | | |
|--|---------------------|----------------------------------|----------------|-----------|---------|
| | | | Total | Irrigated | Rainfed |
| | 1 | Orange | 0.780 | - | 0.780 |
| | 2 | Banana | 0.350 | - | 0.350 |
| | 3 | Pineapple | 0.235 | - | 0.235 |
| | Others (specify) | - | - | - | - |
| | | Horticulture crops Vegetables | Total | Irrigated | Rainfed |
| | 1 | Cabbage | 0.074 | 0.074 | - |
| | 2 | Tomato | 0.049 | 0.049 | - |
| | 3 | Okra | 0.041 | - | 0.041 |
| | 4 | Brinjal | 0.039 | - | 0.039 |
| | 5 | Chilly | 0.154 | - | 0.154 |
| | Others (specify) | - | - | - | - |
| | | Medicinal and Aromatic crops | Total | Irrigated | Rainfed |
| | 1 | Large Cardamom | 0.055 | - | 0.055 |
| | Others (specify) | - | - | - | - |
| | | Plantation crops | Total | Irrigated | Rainfed |
| | 1 | - | - | - | - |

| | | | | | |
|--|---------------------|------------------------------------|--------------|------------------|----------------|
| | Others (Specify) | - | - | - | - |
| | | Fodder crops | Total | Irrigated | Rainfed |
| | 1 | - | - | - | - |
| | Others (Specify) | Grazing land, reserve areas etc | 0.331 | - | 0.331 |
| | | Sericulture | | - | |
| | | Eri | 0.90 | | 0.90 |
| | | Muga | 0.56 | | 0.56 |
| | | Mulberry | 0.4 | | 0.4 |
| | Others (specify) | | | | |

(Source: Statistical Abstract, Arunachal Pradesh, 2007)

| 1.8 | Livestock (Source: Live stock census 2007) | Male ('000) | Female ('000) | Total ('000) |
|------------|---|---------------------|----------------------------------|---------------------|
| | Indigenous cattle | 35.60 | 38.50 | 74.10 |
| | Improved / Crossbred cattle | 1.80 | 2.12 | 3.91 |
| | Buffaloes (local low yielding) | 0.350 | 0.324 | 0.674 |
| | Improved Buffaloes | - | - | - |
| | Goat | 9.52 | 13.99 | 23.51 |
| | Sheep | - | - | 0.043 |
| | Pig | 18.93 | 16.22 | 35.15 |
| | Mithun | - | - | 19.796 |
| | Yak | - | - | - |
| | Others (Horse, mule, donkey etc., specify) | - | - | - |
| | Commercial dairy farms (Number) | - | - | - |
| 1.9 | Poultry | No. of farms | Total No. of birds ('000) | |

| | | | |
|-------------|---|-------------------------------|-------------------------------|
| | Commercial | - | - |
| | Backyard | - | 148.64 |
| 1.10 | Fisheries (Data source: Chief Planning Officer) | | |
| | A. Capture | | |
| | i) Marine (Data Source: Fisheries Department) | No. of fishermen | Boats |
| | | | Mechanized |
| | ii) Inland (Data Source: Fisheries Department) | No. Farmer owned ponds | Reservoirs |
| | | | |
| | B. Culture | | |
| | | Water Spread Area (ha) | Yield (t/ha) |
| | i) Brackish water (Data Source: MPEDA/ Fisheries Department) | | |
| | ii) Fresh water (Data Source: Fisheries Department) | 300 | - |
| | Others (Pond/Farm) | 97.6 | 0.70 |
| | | | Production ('000 tons) |
| | | | 0.180 |
| | | | 0.047 |

(Source: Department of District Veterinary Office; Fishery Department, East Siang, Arunachal Pradesh, 2007.)

1.11 Production and Productivity of major crops (2003-07)

| 1.11 | Name of crop | Kharif | | Rabi | | Summer | | Total | | Crop residue as fodder ('000 tons) |
|--|--------------|---------------------|----------------------|---------------------|----------------------|---------------------|----------------------|---------------------|----------------------|------------------------------------|
| | | Production ('000 t) | Productivity (kg/ha) | Production ('000 t) | Productivity (kg/ha) | Production ('000 t) | Productivity (kg/ha) | Production ('000 t) | Productivity (kg/ha) | |
| Major Field crops (Crops to be identified based on total acreage) | | | | | | | | | | |
| Crop 1 | Paddy | 29.594 | 2252.72 | - | - | - | - | 29.594 | 2252.72 | - |
| Crop 2 | Maize | - | - | - | - | 6.444 | 2235.17 | 6.444 | 2235.17 | - |

| | | | | | | | | | | |
|--|-----------|-------|---------|--------|---------|-------|---------|--------|---------|---|
| Crop 3 | Millet | - | - | - | - | 2.740 | 1234.23 | 2.740 | 1234.23 | - |
| Crop 4 | Pulse | 0.957 | 1021.34 | - | - | - | - | 0.957 | 1021.34 | - |
| Crop 5 | Oilseed | - | - | 1.426 | 821.90 | - | - | 1.426 | 821.90 | - |
| Crop 6 | Potato | - | - | 1.701 | 5785.71 | - | - | 1.701 | 5785.71 | - |
| Crop 7 | Ginger | - | - | - | - | 1.636 | 9242.94 | 1.636 | 9242.94 | - |
| Others | - | - | - | - | - | - | - | - | - | - |
| Major Horticultural crops (Crops to be identified based on total acreage) | | | | | | | | | | |
| Crop 1 | Orange | - | - | - | - | - | - | 7.019 | 1531 | - |
| Crop 2 | Banana | - | - | - | - | - | - | 3.971 | 6894 | - |
| Crop 3 | Pineapple | - | - | - | - | - | - | 3.922 | 3298.57 | - |
| Crop 4 | Cabbage | - | - | 10.240 | 16000 | - | - | 10.240 | 16000 | - |
| Crop 5 | Tomato | - | - | 1.706 | 1333.2 | - | - | 1.706 | 1333.2 | - |
| Crop 6 | Okra | - | - | - | - | 0.054 | 1600 | 0.054 | 1600 | - |
| Crop 7 | Brinjal | - | - | - | - | 0.833 | 2500 | 0.833 | 2500 | - |
| Crop 8 | Chilli | - | - | - | - | 0.238 | 1545.45 | 0.238 | 1545.45 | - |
| Others | - | - | - | - | - | - | - | - | - | - |

(Source: Statistical Abstract, Arunachal Pradesh, 2007)

| 1.12 | Sowing window for 5 major field crops (start and end of normal sowing period) | Paddy | Maize | Oilseed | Ginger | Potato |
|------|--|---|-------|--|--------|---|
| | Kharif- Rainfed | 4 th week June–2 nd week July | - | - | - | - |
| | Kharif-Irrigated | - | - | - | - | - |
| | Rabi- Rainfed | - | - | 1 st week Sept -3 rd week Sept | - | 1 st week Sept-4 th week Sept |
| | Rabi-Irrigated | - | - | - | - | - |


| | | | | | | |
|--|------------------|---|---------------------------------|---|----------------------------------|---|
| | Summer-irrigated | - | - | - | - | - |
| | Summer-rainfed | - | 4th week Mar- 2nd week April | - | 4th week March - 1st week May | - |

| 1.13 | <i>What is the major contingency the district is prone to? (Tick mark)</i> | Regular* | Occasional | None |
|------|--|---|-------------------|-------------|
| | <i>Drought</i> | | √ | |
| | <i>Flood</i> | | √ | |
| | <i>Cyclone</i> | | - | |
| | <i>Hail storm</i> | | - | |
| | <i>Heat wave</i> | | - | |
| | <i>Cold wave</i> | | √ | |
| | <i>Frost</i> | | - | |
| | <i>Sea water intrusion</i> | | - | |
| | <i>Snowfall</i> | | - | |
| | <i>Landslides</i> | | √ | |
| | <i>Earthquake</i> | | √ | |
| | <i>Pests and disease outbreak (specify)</i> | Rice gundhi bug, Citrus stem borer, Tomato fruit borer, Tomato and Potato blights, Bacterial wilt in tomato, Banana pseudo stem borer, Ginger rhizome rot, Rodents in Paddy | - | |
| | <i>Others (like fog, cloud bursting etc.)</i> | | √ | |

*When contingency occurs in six out of 10 years

| | | | |
|------|---|---|---------------|
| 1.14 | Include Digital maps of the district for | Location map of district within State as Annexure I | Enclosed: Yes |
| | | Mean annual rainfall as Annexure 2 | Enclosed: Yes |
| | | Soil map as Annexure 3 | Enclosed: Yes |

Location map of East Siang




Annexure-I: Location Map of District

Annexure-II: Mean Annual Rainfall

| Month | Mean Rain fall (mm) |
|------------------------------|----------------------------|
| January | 365.5 |
| February | 612.4 |
| March | 748.0 |
| April | 315.5 |
| May | 177.6 |
| June | 142.1 |
| July | 22.4 |
| August | 52.2 |
| September | 147.9 |
| October | 42.7 |
| November | 233.8 |
| December | 130.5 |
| Total Annual Rainfall | 3733.6 |

Annexure-III: Soil Map of District



2.0 Strategies for weather related contingencies

2. Drought

2.1 Drought (Rainfed situation)

Drought-Pre-Monsoon (Last week of March to First week of April) Normal

| Condition | Major Farming situation | Normal Crop / Cropping system ^b | Suggested Contingency measures | | |
|--------------------------------------|---|--|--|---|--------------------------------------|
| | | | Change in crop /cropping system including variety | Agronomic measures | Remarks on Implementation |
| Early season drought (delayed onset) | Very gently sloping plains (medium lands) with deep loamy soils | Maize | No change <ul style="list-style-type: none"> Short duration crops/varieties like RCM-1-75, RCM-1-76 Maize + groundnut/soy a bean/rice bean inter cropping. | <ul style="list-style-type: none"> Conservation of pre-monsoon soil moisture through soil/straw/grass mulching practices Hydropriming/ seed soaking in water for 24hr and followed by shade drying before sowing. Application of organic manure before sowing. | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Millets (finger/fox tail millet) | Short duration crops/varieties of finger millet (VR-708, GPU-67), foxtail millet (SR-16, Meera) | | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Pulses (Black gram) | No Change | <ul style="list-style-type: none"> Conservation of pre-monsoon soil moisture through soil/straw/grass mulching practices Application of organic manure before sowing. Hydropriming/ seed soaking in water for 24hr and followed by shade drying before sowing. Grow short duration Black gram varieties USJD 113 and KU 301 | Schemes from Line Deptt. /RKVY/ ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | Maize | No change <ul style="list-style-type: none"> Short duration crops/varieties | <ul style="list-style-type: none"> Conservation of pre-monsoon soil moisture through soil/straw/grass mulching practices Hydropriming/ seed soaking in water for 24hr | Schemes from Line Deptt. /RKVY/ |

| | | | | | |
|--|--|----------------------------------|---|---|--------------------------------------|
| | | | like RCM-1-75, RCM-1-76, Allrounder, HQPM-1, DA-61 A ▪ Maize + groundnut/soya bean/rice bean inter cropping. | and followed by shade drying before sowing. ▪ Application of organic manure before sowing. | ATMA |
| | | Millets (finger/fox tail millet) | No Change Short duration crops/varieties of finger millet (VR-708, GPU-67), foxtail millet (SR-16, Meera) | | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Pulses (Black gram) | No Change | ▪ Conservation of pre-monsoon soil moisture through soil/straw/grass mulching practices ▪ Application of organic manure before sowing. ▪ Hydropriming/ seed soaking in water for 24hr and followed by shade drying before sowing. ▪ Grow short duration Black gram varieties USJD 113 and KU 301 | Schemes from Line Deptt. /RKVY/ ATMA |

Normal onset of pre- monsoon

| Condition | Major Farming situation | Normal Crop/cropping system | Suggested Contingency measures | | |
|--------------------------------------|---|-----------------------------|---|--|---------------------------|
| | | | Crop management | Soil nutrient & moisture conservation measures | Remarks on Implementation |
| Early season drought (Normal onset) | | | | | |
| Normal onset followed by 15-20 | Very gently sloping plains (medium lands) | Maize | ▪ If the germination is less than 30% of optimum plant population, re sowing should be done | ▪ Provide irrigation from the available sources ▪ Mulching with locally available | Schemes from Line Deptt. |

| | | | | | |
|--|---|---------------------------------|---|---|-------------------------------------|
| days dry spell after sowing leading to poor germination/crop stand etc. | with deep loamy soils | | <ul style="list-style-type: none"> ▪ Gap filling to be done to maintain optimum plant density ▪ Foliar application of 1% MOP | material | /RKVY/ATMA |
| | | Millet (finger/fox tail millet) | <ul style="list-style-type: none"> ▪ If the germination is less than 30% of optimum plant population re sowing should be done ▪ Gap filling to be done to maintain optimum plant density ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ATMA |
| | | Pulses (Black gram) | <ul style="list-style-type: none"> ▪ If the germination is less than 30% of optimum plant population re sowing should be done ▪ Gap filling to be done to maintain optimum plant density ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | Maize | <ul style="list-style-type: none"> ▪ If the germination is less than 30% of optimum plant population, re sowing should be done ▪ Gap filling to be done to maintain optimum plant density ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ATMA |
| | | Millet (finger/fox tail millet) | <ul style="list-style-type: none"> ▪ If the germination is less than 30% of optimum plant population re sowing should be done ▪ Gap filling to be done to maintain optimum plant density ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ATMA |
| | | Pulses (Black gram) | <ul style="list-style-type: none"> ▪ If the germination is less than 30% of optimum plant population re sowing should be done ▪ Gap filling to be done to maintain optimum plant density ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ATMA |

| Condition | Major Farming situation | Normal Crop /cropping system | Suggested Contingency measures | | |
|--|---|---------------------------------|---|---|--------------------------------------|
| | | | Crop management | Soil nutrient & moisture conservation measures | Remarks on Implementation |
| Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period) | | | | | |
| Vegetative stage | Very gently sloping plains (medium lands) with deep loamy soils | Maize | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Millet (finger/fox tail millet) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Pulses (Black gram) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | Maize | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Millet (finger/fox tail millet) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Pulses (Black gram) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |

| Condition | Major Farming situation | Normal Crop /cropping system | Suggested Contingency measures | | |
|--|---|---------------------------------|---|---|--------------------------------------|
| | | | Crop management | Soil nutrient & moisture conservation measures | Remarks on Implementation |
| Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period) | | | | | |
| Reproductive stage | Very gently sloping plains (medium lands) with deep loamy soils | Maize | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Millet (finger/fox tail millet) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Pulses (Black gram) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | Maize | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Millet (finger/fox tail millet) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |
| | | Pulses (Black gram) | <ul style="list-style-type: none"> ▪ Weeding ▪ Interculture ▪ Foliar application of 1% MOP | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources ▪ Mulching with locally available material | Schemes from Line Deptt. /RKVY/ ATMA |

| Condition | Major Farming situation | Normal Crop/cropping system | Suggested Contingency measures | | |
|--|---|---------------------------------|-------------------------------------|---|------------------------------------|
| | | | Crop management | Kharif Crop planning | Remarks on Implementation |
| Terminal drought (Early withdrawal of monsoon) | Very gently sloping plains (medium lands) with deep loamy soils | Maize | ▪ Harvest at physiological maturity | ▪ Planning for nursery sowing of Paddy. ▪ Preparation of fields for Sesamum, Soybean | Schemes from Line Deptt./RKVY/ATMA |
| | | Millet (finger/fox tail millet) | ▪ Harvest at physiological maturity | ▪ Planning for nursery sowing of Paddy. ▪ Preparation of fields for Sesamum, Soybean | Schemes from Line Deptt./RKVY/ATMA |
| | | Pulses (Black gram) | ▪ Harvest at physiological maturity | ▪ Planning for nursery sowing of Paddy. ▪ Preparation of fields for Sesamum, Soybean | Schemes from Line Deptt./RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | Maize | ▪ Harvest at physiological maturity | ▪ Planning for nursery sowing of Paddy. ▪ Preparation of fields for Sesamum, Soybean | Schemes from Line Deptt./RKVY/ATMA |
| | | Millet (finger/fox tail millet) | ▪ Harvest at physiological maturity | ▪ Planning for nursery sowing of Paddy. ▪ Preparation of fields for Sesamum, Soybean | Schemes from Line Deptt./RKVY/ATMA |
| | | Pulses (Black gram) | ▪ Harvest at physiological maturity | ▪ Planning for nursery sowing of Paddy. ▪ Preparation of fields for Sesamum, Soybean | Schemes from Line Deptt./RKVY/ATMA |
| | | | | | |

Note : Generally the delay in onset of monsoon by 4 weeks is not applicable

2.2 Drought-Normal onset of Monsoon (1st week of June) Normal

| Condition | Major Farming situation | Normal Crop / Cropping system | Suggested Contingency measures | | |
|--------------------------------------|---|-------------------------------|--|---|-------------------------------------|
| | | | Change in crop /cropping system including variety | Agronomic measures | Remarks on Implementation |
| Early season drought (delayed onset) | Very gently sloping plains (medium lands) with deep loamy soils | WRC/TRC (Paddy) | No change <ul style="list-style-type: none"> Rice vars. RCM-9, RCM-10, RCM 11, CAU-R-1, TTB-404, TTB-303, Mulagavaru, Kanaklata. | <ul style="list-style-type: none"> Spacing of 20x10 cm and 3-5 seedlings/hill Weeding is to be done 30 and 45 days after transplanting. | Schemes from Line Deptt. /RKVY/ATMA |
| | | Soybean | <ul style="list-style-type: none"> Short duration variety JS 335 | <ul style="list-style-type: none"> Weeding is to be done 15 and 30 days after sowing. | Schemes from Line Deptt. /RKVY/ATMA |
| | | Sesame | No Change <ul style="list-style-type: none"> Short duration variety AST-1, St 1683, Gouri, Vinayak | <ul style="list-style-type: none"> Weeding is to be done 15 and 30 days after sowing. | Schemes from Line Deptt. /RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | WRC/TRC (Paddy) | No change <ul style="list-style-type: none"> Rice vars. CAU R-1, CAU R-3, Megha Rice 1 and Megha Rice 2, | <ul style="list-style-type: none"> Spacing of 20x10 cm and 3-5 seedlings/hill Weeding is to be done 30 and 45 days after transplanting. | Schemes from Line Deptt. /RKVY/ATMA |

| Condition | Major Farming situation | Normal Crop / Cropping system | Suggested Contingency measures | | |
|-------------------------------------|---|-------------------------------|---|---|-------------------------------------|
| | | | Crop Management | Agronomic measures | Remarks on Implementation |
| Early season drought (Normal onset) | Very gently sloping plains (medium lands) with deep loamy soils | WRC/TRC (Paddy) | <ul style="list-style-type: none"> Gap filling Weeding to be done Foliar application of 1% MOP Application of organic | <ul style="list-style-type: none"> Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |

| | | | | | |
|------------|--|---------|--|--|-------------------------------------|
| stand etc. | | | manure, wherever possible | | |
| | | Soybean | <ul style="list-style-type: none"> ▪ Timely plant protection of measures for caseworm and leaf folder | | |
| | | | <ul style="list-style-type: none"> ▪ Gap filling ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Application of organic manure, wherever possible ▪ Timely plant protection of measures for leaf roller | <ul style="list-style-type: none"> ▪ Short duration variety JS 335 ▪ Weeding is to be done 15 and 30 days after sowing. ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | | | <ul style="list-style-type: none"> ▪ Gap filling ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Application of organic manure, wherever possible ▪ Timely plant protection of measures for leaf roller | <ul style="list-style-type: none"> ▪ Short duration variety AST-1, St 1683, Gouri, Vinayak ▪ Weeding is to be done 15 and 30 days after sowing. ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | Paddy | <ul style="list-style-type: none"> ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Application of organic manure, wherever possible ▪ Timely plant protection of measures for caseworm and leaf folder | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |

| Condition | Major Farming situation | Normal Crop /cropping system | Suggested Contingency measures | | |
|--|---|------------------------------|---|---|-------------------------------------|
| | | | Crop management | Soil nutrient & moisture conservation measures | Remarks on Implementation |
| Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period) | | | | | |
| Vegetative stage | Very gently sloping plains (medium lands) with deep loamy | WRC/TRC (Paddy) | <ul style="list-style-type: none"> ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for caseworm and leaf | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |

| | | | | | |
|--|---|-----------------|---|---|-------------------------------------|
| | soils | | folder | | |
| | | Soybean | <ul style="list-style-type: none"> ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for leaf roller | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | | Sesames | <ul style="list-style-type: none"> ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for leaf roller | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | WRC/TRC (Paddy) | <ul style="list-style-type: none"> ▪ Weeding to be done ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for Sheath blight, caseworm, leaf folder and stem borer | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |

| Condition | | | Suggested Contingency measures | | |
|---|--|-------------------------------------|---|---|-------------------------------------|
| Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm)period) | Major Farming situation | Normal Crop /cropping system | Crop management | Soil nutrient & moisture conservation measures | Remarks on Implementation |
| Reproductive stage | Very gently sloping plains (medium lands) with deep loamy soils | WRC/TRC (Paddy) | <ul style="list-style-type: none"> ▪ Foliar application of 1% MOP ▪ Timely plant protection measures for gundhi bug, | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | | Soybean | <ul style="list-style-type: none"> ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for pod borer | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | | Sesames | <ul style="list-style-type: none"> ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for capsule borer | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | WRC/TRC (Paddy) | <ul style="list-style-type: none"> ▪ Foliar application of 1% MOP ▪ Timely plant protection of measures for gundhi bug | <ul style="list-style-type: none"> ▪ Provide irrigation from the available sources | Schemes from Line Deptt. /RKVY/ATMA |

| Condition | | | Suggested Contingency measures | | |
|--|--|------------------------------------|-------------------------------------|--|------------------------------------|
| Terminal drought (Early withdrawal of monsoon) | Major Farming situation | Normal Crop/cropping system | Crop management | Rabi Crop planning | Remarks on Implementation |
| | Very gently sloping plains (medium lands) with deep loamy soils | WRC/TRC (Paddy) | ▪ Harvest at physiological maturity | ▪ Planning for zero tillage cultivation of Toria, Pea, Niger etc. ▪ Preparation for cole and solanecous crops | Schemes from Line Deptt./RKVY/ATMA |
| | | Soybean | ▪ Harvest at physiological maturity | ▪ Planning for zero tillage cultivation of Toria, Pea, Niger etc. ▪ Preparation for cole and solanecous crops | Schemes from Line Deptt./RKVY/ATMA |
| | | Sesames | ▪ Harvest at physiological maturity | ▪ Planning for zero tillage cultivation of Toria, Pea, Niger etc. ▪ Preparation for cole and solanecous crops | Schemes from Line Deptt./RKVY/ATMA |
| | Nearly level plains (lowlands) with deep loamy soils | WRC/TRC (Paddy) | ▪ Harvest at physiological maturity | ▪ Planning for zero tillage cultivation of Toria, Pea, Niger etc. ▪ Preparation for cole and solanecous crops | Schemes from Line Deptt./RKVY/ATMA |

2.1.2 Drought-irrigated situation : NA in this district.

2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigation situation)

| Condition | Suggested contingency measure | | | |
|---|---|--|---|--|
| | Vegetative stage | Flowering stage | Crop maturity stage | Post harvest |
| Continuous high rainfall in a short span leading to water logging paddy | Drainage of excess water from the field | Immediate provision of drainage system | ▪ Drain out excess water ▪ Harvest at physiological maturity | ▪ Shifting to a safer place ▪ Dry in shade and in well ventilated space |

| | | | | |
|---------------------|--|--|--|--|
| Maize | Provide drainage | Provide drainage | <ul style="list-style-type: none"> ▪ Drain out excess water ▪ Harvest at physiological maturity | <ul style="list-style-type: none"> ▪ Shifting to a safer place ▪ Dry in shade and in well ventilated space |
| Milllet | Drainage of excess water | Immediate provision of drainage system | <ul style="list-style-type: none"> ▪ Drain out excess water ▪ Harvest at physiological maturity | Proper drying |
| Horticulture | | | | |
| Orange | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ In steep slopes, prepare half moon terraces to prevent soil erosion and leaching loss ▪ If there is physical damage, pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection. ▪ Proper nutrient management to be followed. | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Foliar application of micronutrient/multiplex @ 0.2% should be done to prevent flower drop ▪ Control aphids and mealy bugs etc | <ul style="list-style-type: none"> ▪ If there is physical damage, pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection ▪ Harvesting can be delayed upto 60-75 days by spraying pre-harvest chemical i.e. 2-4D at 20ppm + GA at 10ppm + 0.2% KCl on maturing fruits. ▪ Harvesting can be delayed. In citrus even after full maturity, the fruits can be left on the tree for 2-3 weeks without deterioration which facilitates prolong harvesting. ▪ While picking, the stem end should be cut close to the fruit without damaging the rind. Hence avoiding fungal infection. ▪ Collect the good fruits and store them. Damaged fallen fruits to be disposed off | <ul style="list-style-type: none"> ▪ Fruits are to be stored in well aerated farm shed or house to avoid loses. ▪ Storing at 8 – 10 0 C with 85 – 90 % RH is preferred. |
| Apple | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ In steep slopes, prepare half moon terraces to prevent soil erosion and leaching loss ▪ If there is physical damage, pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection ▪ Nutrient management to be done | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Half moon terraces to be done to prevent nutrient loss ▪ Pruning of damaged brances and application of Bordeaux Paste to be done ▪ Nutrient management along with foliar application micronutrient to be done | <ul style="list-style-type: none"> ▪ Spray 2,4,5-T @ 20ppm or 2,4,5-TCPA @ 15ppm to inhibit fruit drop ▪ Collect the good fruits and store them. Damaged fallen fruits to be separated and disposed off ▪ Necessary to maintain adequate drainage | <ul style="list-style-type: none"> ▪ Stored the fruits for 4-8 months at -1.1 to 0°C and 85-90 % RH. ▪ Spray growth regulators Like Alar @ 1000 ppm to improve storability |

| | | | | |
|----------------|---|---|---|---|
| Pineapple | <ul style="list-style-type: none"> ▪ Make trenches/furrows in between ridges to facilitate drainage of excess water ▪ Remove the excess suckers to maintain the quality of plant ▪ Nutrient management to be followed | <ul style="list-style-type: none"> ▪ Application of Ethephon 2mg in 100-140mg, Bentonite or NAA @ 25ppm or 2, 4-D @ 5-10 ppm should be applied for uniform flower induction. | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Spraying of insecticides and fungicide ▪ Fruits can be protected with locally available material to protect the mature fruit from unusual rains | <ul style="list-style-type: none"> ▪ Store fruits in well aerated farm shed or house to avoid loses. ▪ Pineapples can be stored at a temperature of 7.5-12°C and RH 70-90% for 4 weeks. |
| Kiwifruit | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ In steep slopes, prepare half moon terraces to prevent soil erosion and leaching loss ▪ If there is physical damage, pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection ▪ Nutrient management to be done | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Half moon terraces to be done to prevent nutrient loss ▪ Pruning of damaged branches and application of Bordeaux Paste to be done ▪ Nutrient management along with foliar application micronutrient to be done | <ul style="list-style-type: none"> ▪ Heavy pruning should not done as the fruit will be affected by rain ▪ Drain out excess water | <ul style="list-style-type: none"> ▪ Stored the fruits at 0 to 4°C and 80-90 % RH. ▪ Spray growth regulators Like Alar @ 1000 ppm to improve storability |
| Banana | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping or staking should be done ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done along with application of micronutrient ▪ Propping or staking should be done ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping to be done ▪ Bagging to be done to protect the bunch from unusual rains. ▪ Denavelling to be done to improve the bunch weight (removal of male bud) | <ul style="list-style-type: none"> ▪ Store the fruits/ bunch in well aerated farm shed or house to avoid loses. ▪ Storing at 10 – 12° C with 70 – 80 % RH |
| Large cardamom | <ul style="list-style-type: none"> ▪ It grows luxuriantly in moist and humid climate. So continuous rain is not a problem during its vegetative growth. ▪ Provide adequate drainage ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Rain during flowering is detrimental. So water logging should be avoided. ▪ Proper drainage system should be followed. ▪ Shade regulation may be taken up providing 50-60% shade. | <ul style="list-style-type: none"> ▪ Harvesting can be delayed ▪ Proper drainage system should be followed. | <ul style="list-style-type: none"> ▪ Collect and dry the produce in fuel kiln overnight at 50°-60°C or in drier for 14-18 hours at 45°-50°C |
| Ginger | <ul style="list-style-type: none"> ▪ Provide proper drainage channels to avoid stagnation of water ▪ Earthing up to be done at proper soil moisture level ▪ Nutrient management to be followed ▪ Field bunding to prevent entry of water from surrounding areas. | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up should be followed by manuring. ▪ Field bunding to prevent entry of water from surrounding areas. | <ul style="list-style-type: none"> ▪ Dry weather before harvesting is necessary. So harvesting can be delayed. | <ul style="list-style-type: none"> ▪ Shifting of the produce to a drier place. ▪ Drying to remove excess moisture of produce. |

| | | | | |
|---|--|--|---|--|
| | <ul style="list-style-type: none"> ▪ Spraying of insecticides and fungicide | | | |
| Turmeric | <ul style="list-style-type: none"> ▪ Provide proper drainage channels to avoid stagnation of water ▪ Earthing up to be done at proper soil moisture level ▪ Nutrient management to be followed ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up should be followed by manuring. ▪ Field bunding to prevent entry of water from surrounding areas. | <ul style="list-style-type: none"> ▪ Dry weather before harvesting is necessary. So harvesting can be delayed. | <ul style="list-style-type: none"> ▪ Shifting of the produce to a drier place. ▪ Drying to remove excess moisture of produce. |
| Vegetables (cucurbits) | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up to be done at proper soil moisture condition followed by manuring ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Staking should be properly followed. Rainy season crops can be trained on a bower made of bamboos and sticks. | <ul style="list-style-type: none"> ▪ Spray maleic hydrazine (MH) and 2, 4-5 tri-iodobenzoic acid (TIBA) @ 50ppm for Sex expression. Boron @ 3ppm and calcium @ 20ppm is also effective. ▪ Provision of drainage to remove excess water. ▪ Earthing up followed by manuring ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Take up proper plant protection measures | <ul style="list-style-type: none"> ▪ Fruits to be harvested immediately without causing injury to fruits ▪ Remove all damaged fruit ▪ Take up appropriate plant protection measures | <ul style="list-style-type: none"> ▪ The fruits can be stored for 2-3 weeks at 15-20°C and RH 75% in a well-ventilated chamber |
| Heavy rainfall with high speed winds in a short span | | | | |
| Horticulture | | | | |
| Orange | <ul style="list-style-type: none"> ▪ Earthing up of young plants to avoid uprooting due to wind. ▪ Provide proper drainage facilities. ▪ Staking to avoid falling off of plants ▪ In steep slopes, prepare half moon terraces to prevent soil erosion and leaching loss ▪ Pruning of damage branches and | <ul style="list-style-type: none"> ▪ Wind break around the orchard to protect crop from wind damage ▪ Provide proper drainage ▪ Nutrient management to be followed along with foliar spray of micronutrient ▪ Pruning of damage branches | <ul style="list-style-type: none"> ▪ Propping heavy bearing tree and weak tree by bamboo pole. ▪ Harvesting can be delayed upto 60-75 days by spraying pre-harvest chemical i.e. 2-4D at 20ppm + GA at 10ppm + 0.2% Kcl on maturing fruits. ▪ Pruning of damage branches | <ul style="list-style-type: none"> ▪ Fruits are to be stored in well aerated farm shed or house to avoid loses. ▪ Pack the fruit in perforated polythene bag, boxes, crates, etc. and store at temperature of 10-11°C & 92 % RH. |

| | | | | |
|-----------|--|---|--|---|
| | <p>application of Bordeaux paste should be done to prevent secondary infection</p> <ul style="list-style-type: none"> Proper nutrient management to be followed | <p>and application of Bordeaux paste should be done to prevent secondary infection</p> | <p>and application of Bordeaux paste should be done to prevent secondary infection</p> | |
| Apple | <ul style="list-style-type: none"> Earthing up of young plants to avoid uprooting due to wind. Provide proper drainage facilities. Staking to be done to avoid falling off of plants. In steep slopes, prepare half moon terraces to prevent soil erosion and leaching loss Pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection Proper nutrient management to be followed | <ul style="list-style-type: none"> Provision of drainage to remove excess water. Wind break around the orchard Maintain the half moon terraces to avoid soil nutrient loss Proper nutrient management to be followed along with foliar application of micronutrient Prune out all damage branches with appropriate plant protection measures | <ul style="list-style-type: none"> Harvest ripe fruits Propping heavy bearing tree and weak tree by bamboo pole. Use of plant bio-regulators to delay ripening with Daminozide or Alar @ 1000ppm sprayed before 60 days before harvest. | <ul style="list-style-type: none"> Store fruits for 4-8 months at -1.1 to 0°C and 85-90 % RH. |
| Pineapple | <ul style="list-style-type: none"> Earthing up plants for better development and anchorage. Make trenches/furrows in between ridges to facilitate drainage of excess water. Nutrient management to be followed | <ul style="list-style-type: none"> Earthing up to prevent uprooting. Provide proper drainage Nutrient management to be followed Spray NAA @ 25ppm or 2, 4-D @ 5-10 ppm should be applied for uniform flower induction. | <ul style="list-style-type: none"> Fruits can be protected with locally available material to protect the mature fruit from unusual rains Spraying of insecticides and fungicide Earthing up plants for better development and anchorage. Make trenches/furrows in between ridges to facilitate drainage of excess water | <ul style="list-style-type: none"> Store fruits in well aerated farm shed or house to avoid loses. Pineapples can be stored at a temperature of 7.5-12°C and RH 70-90% for 4 weeks. |
| Kiwifruit | <ul style="list-style-type: none"> Provide proper drainage Support the plant using T-Bar system In steep slopes, prepare half moon terraces to prevent soil erosion and leaching loss If there is physical damage, pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection Nutrient management to be done | <ul style="list-style-type: none"> Provide proper drainage Half moon terraces to be done to prevent nutrient loss Pruning of damaged branches and application of Bordeaux Paste to be done Nutrient management along with foliar application micronutrient to be done | <ul style="list-style-type: none"> Heavy pruning should not done as the fruit will be affected by rain Drain out excess water Maintain the plant using T-Bar trellis supporting system Nutrient management along with foliar application micronutrient to be done | <ul style="list-style-type: none"> Stored the fruits at 0 to 4°C and 80-90 % RH. Spray growth regulators Like Alar @ 1000 ppm to improve storability |

| | | | | |
|------------------------|--|---|---|---|
| Banana | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping or staking should be done ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done along with application of micronutrient ▪ Propping or staking should be done ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping to be done ▪ Bagging to be done to protect the bunch from unusual rains. ▪ Denavelling to be done to improve the bunch weight (removal of male bud) | <ul style="list-style-type: none"> ▪ Store the fruits/ bunch in well aerated farm shed or house to avoid loses. ▪ Storing at 10 – 12° C with 70 – 80 % RH |
| Large cardamom | <ul style="list-style-type: none"> ▪ For newly planted crops, staking should be provided. ▪ Provide adequate drainage ▪ Spraying of insecticides and fungicid ▪ Follow proper nutrient management ▪ Earthing up to be done | <ul style="list-style-type: none"> ▪ Proper drainage system should be followed. ▪ Follow proper nutrient management ▪ Earthing up to prevent uprooting. | <ul style="list-style-type: none"> ▪ Harvest at physiological maturity stage or can be delayed ▪ Proper drainage system should be followed | <ul style="list-style-type: none"> ▪ Collect the harvest and dry the produce in fuel kiln overnight at 50°-60°C or in drier for 14-18 hours at 45°-50°C |
| Ginger | <ul style="list-style-type: none"> ▪ Provide proper drainage channels to avoid stagnation of water ▪ Earthing up to be done at proper soil moisture level ▪ Nutrient management to be followed ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up should be followed by manuring. ▪ Field bunding to prevent entry of water from surrounding areas. | <ul style="list-style-type: none"> ▪ Harvest at physiological maturity stage. | <ul style="list-style-type: none"> ▪ Shifting of the produce to a drier place. ▪ Drying to remove excess moisture of produce (moisture level 10%) |
| Turmeric | <ul style="list-style-type: none"> ▪ Provide proper drainage channels to avoid stagnation of water ▪ Earthing up to be done at proper soil moisture level ▪ Nutrient management to be followed ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up should be followed by manuring. ▪ Field bunding to prevent entry of water from surrounding areas. | <ul style="list-style-type: none"> ▪ Dry weather before harvesting is necessary. So harvesting can be delayed. | <ul style="list-style-type: none"> ▪ Shifting of the produce to a drier place. ▪ Drying to remove excess moisture of produce. |
| Vegetables (cucurbits) | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up to be followed ▪ Ensure proper staking of crop wherever required ▪ Field bunding to prevent entry of water from surrounding areas. | <ul style="list-style-type: none"> ▪ Spray maleic Hydrazide @ 50ppm aqueous solution at 2 and 4 leaf stages to stimulate vine growth, giving more female flowers. ▪ Provision of drainage to remove excess water. | <ul style="list-style-type: none"> ▪ Fruits to be harvested immediately without causing injury to fruits ▪ Remove all damaged fruit ▪ Take up appropriate plant protection measures | <ul style="list-style-type: none"> ▪ The fruits can be stored for 2-3 weeks at 15-20°C and RH 75% in a well-ventilated chamber. |

| | | | | |
|--|--|---|--|---|
| | | <ul style="list-style-type: none"> ▪ Wind break around the orchard to protect crop from wind damage ▪ Earthing up and propping to prevent uprooting. ▪ Field bunding to prevent entry of water from surrounding areas. | | |
| Outbreak of pests and diseases due to unseasonal rains : NA | | | | |
| Paddy (Blast) | <ul style="list-style-type: none"> ▪ Use trap crops for prediction of disease. ▪ Removal and destruction of weed hosts in the field bunds and channels | <ul style="list-style-type: none"> ▪ Spraying of Mancozeb @ 2g/ltr or spraying of Carbendazim @ 1 g/ltr. | <ul style="list-style-type: none"> ▪ Drain out excess water to avoid flooded conditions. | <ul style="list-style-type: none"> ▪ Sun drying to prevent spoilage and sprouting of the harvested grains. |
| Paddy (Brown Spot) | -Do- | -Do- | -Do- | -Do- |
| Paddy (Bacterial leaf blight) | <ul style="list-style-type: none"> ▪ Destruction of weed hosts. | <ul style="list-style-type: none"> ▪ Spraying of streptomycin and tetracycline. | <ul style="list-style-type: none"> ▪ Drain out excess water to avoid flooded conditions. | -Do- |
| Paddy (Yellow Stem Borer) | <ul style="list-style-type: none"> ▪ Collection and destruction of egg masses. | <ul style="list-style-type: none"> ▪ Spraying of Chloropyrifos 20 EC @ 0.02 %. | <ul style="list-style-type: none"> ▪ Harvesting at the right stage. | -Do- |
| Paddy (Gall Midge) | <ul style="list-style-type: none"> ▪ Removal of alternate host plants including weeds and grasses and destruction of infected plants. | <ul style="list-style-type: none"> ▪ Providing proper drainage system. | <ul style="list-style-type: none"> ▪ Harvesting at the right stage. | -Do- |
| Maize (Stalk rot) | <ul style="list-style-type: none"> ▪ Removal of accumulated water around the stalks by proper drainage. | <ul style="list-style-type: none"> ▪ Rouging of affected plant and its destruction. | <ul style="list-style-type: none"> ▪ Spraying of streptocycline @ 0.020 %. | <ul style="list-style-type: none"> ▪ Sun drying of the harvested cob to prevent spoilage. |
| Horticulture | | | | |
| Orange (Citrus Leaf miner) | <ul style="list-style-type: none"> ▪ Spraying of Fenvalerate and Cypermethrin for controlling leaf miner. | <ul style="list-style-type: none"> ▪ Spraying of Fenvalerate and Cypermethrin for controlling leaf miner. | <ul style="list-style-type: none"> ▪ Harvesting at the right stage and proper handling of the produce. | <ul style="list-style-type: none"> ▪ Store in cool place in crates, boxes etc |
| Orange (Citrus butterfly) | <ul style="list-style-type: none"> ▪ Hand picking of caterpillars and pupae in the nursery. | <ul style="list-style-type: none"> ▪ Spraying of Neem formulation to control citrus butterfly. | Do | <ul style="list-style-type: none"> ▪ Store in cool place in crates, boxes etc |
| Orange (Powdery mildew in citrus) | <ul style="list-style-type: none"> ▪ Spraying of wettable sulphur and carbendazim to control powdery mildews. | <ul style="list-style-type: none"> ▪ Spraying of wettable sulphur, bavistin (0.1 %) and calixin (0.1 %). | <ul style="list-style-type: none"> ▪ Spraying of wettable sulphur and carbendazim to control powdery mildews. | <ul style="list-style-type: none"> ▪ Store in cool place in crates, boxes etc. |
| Tomato | <ul style="list-style-type: none"> ▪ Removal of accumulated water by proper drainage. ▪ Destroy the heavily infested/infected | <ul style="list-style-type: none"> ▪ Spraying of Sulfax @ 2 g/ltr of water. | <ul style="list-style-type: none"> ▪ Harvesting at the right stage and proper handling. | <ul style="list-style-type: none"> ▪ Store in cool/dry place packed in crates, boxes etc. |

| | | | | |
|-------------------|--|--|--|--|
| | plant parts. | | | |
| Brinjal | <ul style="list-style-type: none"> ▪ Removal of accumulated water by proper drainage. ▪ Destroy the heavily infested/infected plant parts. | <ul style="list-style-type: none"> ▪ Spraying of Sulfex @ 2 g/lit of water. ▪ Soil dranching with captan/Tiram @ 2/lit of water | <ul style="list-style-type: none"> ▪ Harvesting at the right stage and proper handling of the produce. | <ul style="list-style-type: none"> ▪ Store in cool/dry place packed in crates, boxes etc. |
| Cabbage | <ul style="list-style-type: none"> ▪ Removal of accumulated water by proper drainage. ▪ Destroy the badly infested/infected plant parts. | <ul style="list-style-type: none"> ▪ Spraying of Sulfex @ 2 g/lit of water. ▪ Soil dranching with captan/Tiram. @ 2/lit of water ▪ Streptocycline spray | <ul style="list-style-type: none"> ▪ Harvesting at the right stage and proper handling of the produce. | <ul style="list-style-type: none"> ▪ Store in cool/dry place |
| Cucurbits | <ul style="list-style-type: none"> ▪ Manual collection & destruction of eggs/grubs/larvae. | <ul style="list-style-type: none"> ▪ Spraying of carbaryl against leaf eating caterpillars, Metalaxyl against Powdery mildew, Carbendazim against leaf spot & blight | <ul style="list-style-type: none"> ▪ Spraying of Malathion against fruit fly. | <ul style="list-style-type: none"> ▪ Store in cool/dry place |
| Large Cardamom | <ul style="list-style-type: none"> ▪ Proper drainage. ▪ Uprooting and destruction of Chirke and Foorkey infected cardamom plants. | <ul style="list-style-type: none"> ▪ Removal of affected plant from the field. | <ul style="list-style-type: none"> ▪ Harvesting at the right stage and proper handling of the produce. | <ul style="list-style-type: none"> ▪ Quick drying of harvested capsule. |
| Ginger (Soft rot) | <ul style="list-style-type: none"> ▪ Removal of accumulated water in the field by proper drainage. | <ul style="list-style-type: none"> ▪ Removal and destruction of affected plants. | <ul style="list-style-type: none"> ▪ Spraying with Blitox – 50 (3 g/lit) or Dithane – Z-78 (2.5 g / lit). | <ul style="list-style-type: none"> ▪ Store in cool/dry place |

2.3 Floods

| Condition | Suggested contingency measure | | | |
|---|--|--|---|---|
| | Seedling / nursery stage | Vegetative stage | Reproductive stage | At harvest |
| Transient water logging/partial inundation | | | | |
| Rice | <ul style="list-style-type: none"> ▪ Drainage of the Nursery bed. ▪ Re -sowing if not possible | <ul style="list-style-type: none"> ▪ Drainage of excess water. ▪ Gap filling In partially damaged field by redistributing the tillers. ▪ Management of pests & diseases | <ul style="list-style-type: none"> ▪ Drainage of excess water. If flood comes during reproductive stage, emphasis should be given on forthcoming rabi crops. ▪ Utilization of residual soil moisture and use of recharged soil profile for growing pulses | <ul style="list-style-type: none"> ▪ Drainage of excess water. If flood comes during reproductive stage, emphasis should be given on forthcoming rabi crops. ▪ Utilization of residual soil moisture and use of recharged soil profile for growing pulses |
| Horticulture/Plantation crops | | | | |

| | | | | |
|------------------------|--|--|---|---|
| Banana | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping or staking should be done ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping or staking should be done ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provide proper drainage ▪ Nutrient management to be done ▪ Propping to be done | <ul style="list-style-type: none"> ▪ Store the fruits/ bunch in well aerated farm shed or house to avoid loses. ▪ Storing at 10 – 12° C with 70 – 80 % RH |
| Ginger | <ul style="list-style-type: none"> ▪ Provide proper drainage channels to avoid stagnation of water ▪ Earthing up to be done at proper soil moisture level ▪ Nutrient management to be followed ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up should be followed by manuring. ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Application of fungicide and insecticides | <ul style="list-style-type: none"> ▪ Harvest at physiological maturity stage or can delay harvesting | <ul style="list-style-type: none"> ▪ Shifting of the produce to drier place. |
| Turmeric | <ul style="list-style-type: none"> ▪ Provide proper drainage channels to avoid stagnation of water ▪ Earthing up to be done at proper soil moisture level ▪ Nutrient management to be followed ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Spraying of insecticides and fungicide | <ul style="list-style-type: none"> ▪ Provision of drainage to remove excess water. ▪ Earthing up should be followed by manuring. ▪ Field bunding to prevent entry of water from surrounding areas. ▪ Application of fungicide and insecticides | <ul style="list-style-type: none"> ▪ Harvest at physiological maturity stage or can delay harvesting | <ul style="list-style-type: none"> ▪ Shifting of the produce to drier place |
| Vegetables (cucurbits) | <ul style="list-style-type: none"> ▪ Proper drainage of the nursery bed, If not possible go for re-sowing. ▪ Raised bed method should be followed in the nursery. ▪ Earthing up to be followed ▪ Ensure proper staking of | <ul style="list-style-type: none"> ▪ Proper drainage of the nursery bed, If not possible go for re-sowing. ▪ Earthing up to be followed ▪ Ensure proper staking of crop wherever required ▪ Field bunding to prevent entry of water from surrounding | <ul style="list-style-type: none"> ▪ Drainage of excess water. If flood comes during reproductive stage, emphasis should be given on forthcoming rabi crops ▪ Growing of cole crops or winter vegetables after receding flood water and | <ul style="list-style-type: none"> ▪ Shifting of the produce to drier place and store fruits in a well-ventilated chamber |

| | | | | |
|--|---|--|--|----|
| | crop wherever required ▪ Field bunding to prevent entry of water from surrounding areas. | areas. ▪ Follow appropriate nutrient management practices | adoption of integrated farming system to obtain more income and to compensate the loss during kharif vegetables. | |
| Continuous submergence for more than 2 days² | | | | |
| Crop1 | NA | NA | NA | NA |
| Horticulture / Plantation crops | | | | |
| Crop1 (specify) | NA | NA | NA | NA |
| Sea water intrusion³ | | | | |
| Crop1 | NA | NA | NA | NA |

2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone: Not Applicable

| Extreme event type | Suggested contingency measure ^r | | | |
|------------------------------|--|---|---|------------|
| | Seedling / nursery stage | Vegetative stage | Reproductive stage | At harvest |
| Horticulture | | | | |
| Heat Wave^p | | | | |
| Orange | NA | NA | NA | NA |
| Apple | NA | NA | NA | NA |
| Pineapple | NA | NA | NA | NA |
| Kiwifruit | NA | NA | NA | NA |
| Banana | NA | NA | NA | NA |
| Large Cardamom | NA | NA | NA | NA |
| Ginger | NA | NA | NA | NA |
| Turmeric | NA | NA | NA | NA |
| Horticulture | | | | |
| Cold wave^q | | | | |
| Orange | NA | NA | NA | NA |
| Apple | NA | NA | NA | NA |
| Pineapple | NA | NA | NA | NA |
| Kiwifruit | NA | NA | NA | NA |
| Banana | ▪ Protect the plant by construction of | ▪ Protect the plant by construction of wind | ▪ Protect the plant by construction of wind | NA |

| | | | | |
|---------------------|---|---|---|--|
| | <ul style="list-style-type: none"> wind brakes made of shade net. ▪ Maintain the seedling in polyhouse | brakes made of shade net | <ul style="list-style-type: none"> brakes made of shade net ▪ Protect the bunch by bagging with polyethylene bag or jute bag | |
| Large Cardamom | NA | NA | NA | NA |
| Ginger | NA | NA | NA | NA |
| Turmeric | NA | NA | NA | NA |
| Horticulture | | | | |
| Frost | | | | |
| Orange | NA | NA | NA | NA |
| Apple | NA | NA | NA | NA |
| Pineapple | NA | NA | NA | NA |
| Kiwifruit | NA | NA | NA | NA |
| | <ul style="list-style-type: none"> ▪ Protect the plant by construction of wind brakes made of shade net. ▪ Maintain the seedling in polyhouse | <ul style="list-style-type: none"> ▪ Protect the plant by construction of wind brakes made of shade net | <ul style="list-style-type: none"> ▪ Protect the plant by construction of wind brakes made of shade net ▪ Protect the bunch by bagging with polyethylene bag or jute bag | NA |
| Banana | | | | |
| Large Cardamom | NA | NA | NA | NA |
| Ginger | NA | NA | NA | NA |
| Turmeric | NA | NA | NA | NA |
| Horticulture | | | | |
| Hailstorm | | | | |
| Orange | <ul style="list-style-type: none"> ▪ Nursery raising under polyhouse. | <ul style="list-style-type: none"> ▪ Pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection ▪ Nutrient management to be followed along with foliar spray of micronutrient | <ul style="list-style-type: none"> ▪ Pruning of damage branches and application of Bordeaux paste should be done to prevent secondary infection ▪ Nutrient management to be followed along with foliar spray of micronutrient | <ul style="list-style-type: none"> ▪ Harvest ripe fruit |
| Apple | <ul style="list-style-type: none"> ▪ Nursery raising under polyhouse. | <ul style="list-style-type: none"> ▪ Pruning of damage branches and application of Bordeaux | <ul style="list-style-type: none"> ▪ Pruning of damage branches and application of Bordeaux | <ul style="list-style-type: none"> ▪ Harvest ripe fruit |

| | | | | |
|------------------------|---|--|--|---|
| | | paste should be done to prevent secondary infection ▪ Nutrient management to be followed along with foliar spray of micronutrient | paste should be done to prevent secondary infection ▪ Nutrient management to be followed along with foliar spray of micronutrient | |
| Pineapple | NA | ▪ Shade regulation may be followed | NA | ▪ Harvest and value addition |
| Kiwifruit | ▪ Nursery raising under polyhouse | ▪ Nutrient management to be followed along with foliar spray of micronutrient | ▪ Nutrient management to be followed along with foliar spray of micronutrient | ▪ Harvest ripe fruits |
| Banana | ▪ Nursery raising under polyhouse | ▪ Follow nutrient management | ▪ Bagging the fruit bunch with polyethylene bag or jute bag | ▪ Harvest the mature bunch |
| Large Cardamom | ▪ Nursery raising under polyhouse. | ▪ Shade regulation may be followed by planting trees providing 50-60% shade. Ultis cum large cardamom plantation is highly recommended | NA | NA |
| Ginger | ▪ Nursery raising under polyhouse. | ▪ Shade regulation may be followed | NA | NA |
| Turmeric | ▪ | ▪ | | |
| Vegetables (cucurbits) | ▪ Nursery raising under polyhouse. ▪ Provide shade to protect from damage or resowing of the crops | ▪ Polyhouse cultivation & proper irrigation | ▪ Polyhouse cultivation & proper irrigation ▪ Proper crop management for the succeeding years | ▪ Picking of fruits at right edible stage depends upon individual varieties and marketing requirements. Fruits are harvested, packed in baskets and transported to markets. |
| Horticulture | | | | |
| Cyclone | NA | NA | NA | NA |
| Orange | NA | NA | NA | NA |
| Apple | NA | NA | NA | NA |
| Pineapple | NA | NA | NA | NA |
| Kiwifruit | NA | NA | NA | NA |
| Banana | NA | NA | NA | NA |
| Large Cardamom | NA | NA | NA | NA |

| | | | | |
|---|----|----|----|----|
| Ginger | NA | NA | NA | NA |
| Turmeric | NA | NA | NA | NA |
| Sand deposition or heavy siltation | | | | |
| Specify crop /horticulture/plantation | NA | NA | NA | NA |

2.5 Contingent strategies for Livestock, Poultry & Fisheries

2.5.1 Livestock

| | Suggested contingency measures | | |
|-------------------------------|---|--|--|
| | Before the event ^s | During the event | After the event |
| Drought | | | |
| Feed and fodder availability | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories. ▪ Awareness on fodder cultivation & identification of locally available, natural fodder of area. ▪ Excess fodder may be stored as hay/silage or converted into feed block in the flush season, for lean period. ▪ Stacking of paddy straws. | <ul style="list-style-type: none"> ▪ Use of unconventional feed/fodders resources. ▪ Grazing in the peri peri of forest areas. ▪ Feeding according to body weight requirement ▪ Improvement of the poor quality roughages (urea treatment, soaking, poultry litter(> 37%). ▪ Use of feed additives to improve digestibility. ▪ use of stored Hay and Silage | <ul style="list-style-type: none"> ▪ Avail the benefits of schemes under drought, from state or central for feeds and fodder. ▪ Supplementary feeding of livestock to regain the general physiological imbalanced. ▪ Proper irrigation of fodder plot and cultivation of leguminous fodders to meet the demand of green fodders |
| Drinking water | <ul style="list-style-type: none"> ▪ Construction of water harvesting structures. ▪ Harvesting rain water & water from natural source ▪ Developing watershed areas. | <ul style="list-style-type: none"> ▪ Use of stored water from water harvesting structure. ▪ Fetching water from watershed areas and natural stream/river. ▪ Avail subsidy water supply through tankers from sate or central Govt. | <ul style="list-style-type: none"> ▪ Submitting a memorandum to sate or central Govt. regarding amount of water shortfall during drought and action to be initiate accordingly. ▪ Construction of permanent water harvesting structure with a planning to fulfill the water requirement during drought. |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ selective culling of disease animal ▪ Submitting a memorandum to sate or central Govt. regarding the loss of animal due to Drought and remedies to be taken accordingly for future. |

| | | | |
|-------------------------------|--|---|---|
| | <p>the instruction of Do & Don'ts from experts.</p> <ul style="list-style-type: none"> ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. ▪ Proper ventilation system of Housing to reduce heat stress. | | <ul style="list-style-type: none"> ▪ Mini vaccine unit could be establish for covering a perimeter 30-50 km. |
| Floods | | | |
| Feed and fodder availability | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories. ▪ Awareness on fodder cultivation & identification of locally available, natural fodder of the area. ▪ Excess fodder may be stored as hay/silage or converted into feed block in the flush season, for lean period. ▪ Stacking of paddy straws. ▪ Installation of feed block machines and creating feed/fodder block banks to be used in emergency. | <ul style="list-style-type: none"> ▪ Avoid feeding of damp feeds and fodders ▪ Storage of feeds and fodder in high raised platform. ▪ Use of unconventional feed/fodders resources (water hyacinth) ▪ Shifting of livestock to high raised areas. ▪ Use of feed additives to improve digestibility. ▪ Provision of UMB etc. ▪ Use of stored Hay and Silage | <ul style="list-style-type: none"> ▪ Submitting a reports, damage caused by flood to feed and standing fodder ▪ Supplementary feeding of livestock to regain the general physiological imbalanced. ▪ Proper irrigation of folder plot and cultivation of leguminous fodders to meet the demand of green fodders. ▪ Avail the benefits of schemes under flood, from state or central for feeds and fodder. |
| Drinking water | <ul style="list-style-type: none"> ▪ Storage of safe drinking water in community tanks / water harvesting structures which is not prone to seepage of flood water. ▪ Installation of large sized sand filters with charcoal. ▪ Tying up with PHED Deptt. of neighboring district to supply water at needy time. ▪ Creating awareness amongst public how to conserve water and judiciously use in flood situation. | <ul style="list-style-type: none"> ▪ Chlorination of the drinking water and use of sand filter ▪ Incorporation of aquatic plants in feeds as a supplementary source of water ▪ If possible supply of fresh drinking water from nearby district. | <ul style="list-style-type: none"> ▪ Cleaning of water storage tanks, canals and drainage system. ▪ Cleaning and disinfection of water source with suitable water purifying agent, available in the area as per the recommended dose. ▪ Relief for damaged tanks and community pipe line for reconstruction. ▪ Avoid shallow source of water |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Vaccination of FMD, BQ and HS. ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal |

| | | | |
|-------------------------------|---|---|--|
| | <p>the situation if arise.</p> <ul style="list-style-type: none"> ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. ▪ Construction of shelters in high raised areas. | | <ul style="list-style-type: none"> ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to flood and remedies to be taken accordingly for future. |
| Cyclone | NA | NA | NA |
| Feed and fodder availability | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories. ▪ Proper storage of feeds and fodder in well constructed house ▪ Planting of trees as a wind break in farm area ▪ Excess fodder may be stored as hay/silage or converted into feed block in the flush season, for lean period. ▪ Stacking of paddy straws. | <ul style="list-style-type: none"> ▪ Avoid feeding grazing in open field ▪ Animal should be confined in well construct house. ▪ Use of feed additives to improve digestibility. ▪ Provision of UMB etc. ▪ Use of stored Hay and Silage | <ul style="list-style-type: none"> ▪ Submitting a reports, damage caused by cyclone of standing fodder ▪ Avail the benefits of schemes under flood, from state or central for feeds and fodder. |
| Drinking water | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Storage of safe drinking water in community tanks / water harvesting structures ▪ Creating awareness amongst public how to conserve water and judiciously use in flood situation. ▪ Tying up with PHED Deptt. of neighboring district to supply water at needy time. | <ul style="list-style-type: none"> ▪ Chlorination of the drinking water and use of sand filter ▪ Provide fresh potable water | <ul style="list-style-type: none"> ▪ Cleaning of water storage tanks, canals and drainage system. ▪ Cleaning and disinfection of water source with suitable water purifying agent, available in the area as per the recommended dose. ▪ Relief for damaged tanks and community pipe line for reconstruction. ▪ Avoid shallow source of water |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ selective culling of injured animal | <ul style="list-style-type: none"> ▪ Immediate attention to the ailing animals. ▪ selective culling of injured animal ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Sanitization of the shed and surrounding |

| | | | |
|--------------------------------|---|---|--|
| | <p>the situation if arise.</p> <ul style="list-style-type: none"> ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | | <p>areas.</p> <ul style="list-style-type: none"> ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to flood and remedies to be taken accordingly for future. |
| Heat wave | | | |
| Cattle | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Increase the concentrate feed amount and reduce the roughage diet. ▪ Adlib provision of potable water | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to heat wave ▪ reduce upto 20% of the ration ▪ provide nutretical ▪ Adlib provision of potable water ▪ Avoid movement of animal ▪ Sprinkling of water during the extreme heat to the animal ▪ Breeding should be done in morning hours. | <ul style="list-style-type: none"> ▪ Adlib provision of potable water ▪ Analysis of the present experience and remodeling of housing structure. ▪ provide nutretical |
| Health and disease management | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Ensure livestock insurance ▪ Deworming and vaccination ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication | <ul style="list-style-type: none"> ▪ Life saving treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ Oral supplementation of electrolyte and medicines | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ Selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. |

| | | | |
|--------------------------------|--|---|--|
| | and transportation facilities in every dispensary / clinic for consultations. | | |
| Mithun | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Increase the concentrate feed amount and reduce the roughage diet. ▪ Adlib provision of potable water | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to heat wave ▪ reduce upto 20% of the ration ▪ provide nutretical ▪ Adlib provision of potable water ▪ Avoid movement of animal ▪ Sprinkling of water during the extreme heat to the animal ▪ Breeding should be done in morning hours. | <ul style="list-style-type: none"> ▪ Adlib provision of potable water ▪ Analysis of the present experience and remodeling of housing structure. ▪ provide nutretical |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ selective culling of injured animal | <ul style="list-style-type: none"> ▪ Immediate attention to the ailing animals. ▪ selective culling of injured animal ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Sanitization of the shed and surrounding areas. ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to flood and remedies to be taken accordingly for future. |
| Goat/Sheep | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to heat wave ▪ reduce upto 20% of the ration ▪ provide nutretical ▪ Adlib provision of potable water ▪ Avoid movement of animal ▪ Sprinkling of water during the extreme heat to the animal | <ul style="list-style-type: none"> ▪ Adlib provision of potable water ▪ Analysis of the present experience and remodeling of housing structure. ▪ provide nutretical |

| | | | |
|--------------------------------|--|---|--|
| | <ul style="list-style-type: none"> ▪ Increase the concentrate feed amount and reduce the roughage diet. ▪ Adlib provision of potable water | <ul style="list-style-type: none"> ▪ Breeding should be done in morning hours. | |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ selective culling of injured animal | <ul style="list-style-type: none"> ▪ Immediate attention to the ailing animals. ▪ selective culling of injured animal ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Sanitization of the shed and surrounding areas. ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to flood and remedies to be taken accordingly for future. |
| Pig | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Increase the concentrate feed amount and reduce the roughage diet. ▪ Adlib provision of potable water | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to heat wave ▪ reduce upto 20% of the ration ▪ provide nutretical ▪ Adlib provision of potable water ▪ Avoid movement of animal ▪ Sprinkling of water during the extreme heat to the animal ▪ Breeding should be done in morning hours. | <ul style="list-style-type: none"> ▪ Adlib provision of potable water ▪ Analysis of the present experience and remodeling of housing structure. ▪ provide nutretical |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ selective culling of injured animal | <ul style="list-style-type: none"> ▪ Immediate attention to the ailing animals. ▪ selective culling of injured animal ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Sanitization of the shed and surrounding areas. ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal |

| | | | |
|--------------------------------|--|---|--|
| | <ul style="list-style-type: none"> from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | | due to flood and remedies to be taken accordingly for future. |
| Cold wave | | | |
| Cattle | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Feed balance ration to withstand the cold wave prior to occurrence. | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to cold wave ▪ provide extra bedding materials ▪ feed extra ration along with mineral and vitamin supplements to withstand cold wave ▪ | <ul style="list-style-type: none"> ▪ Analysis of the present experience and remodeling of housing structure. |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. |
| Mithun | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Feed balance ration to withstand the cold wave prior to occurrence. | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to cold wave ▪ provide extra bedding materials ▪ feed extra ration along with mineral and vitamin supplements to withstand cold wave ▪ | <ul style="list-style-type: none"> ▪ Analysis of the present experience and remodeling of housing structure. |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, | <ul style="list-style-type: none"> ▪ 1. Mass awareness cum Health camp and symptomatically prompt treatment accordingly. | <ul style="list-style-type: none"> ▪ 1. Mass awareness cum Health camp and symptomatically prompt treatment accordingly. |

| | | | |
|--------------------------------|--|--|--|
| | <ul style="list-style-type: none"> vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ 2. Supplementary feeding of vitamin and mineral to improve general body health. | <ul style="list-style-type: none"> ▪ 2. Immediate attention to the ailing animals. ▪ 3. Sanitization of the shed and surrounding areas. ▪ 4. selective culling of animal ▪ 5. Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. |
| Pig | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Feed balance ration to withstand the cold wave prior to occurrence. | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to cold wave ▪ provide extra bedding materials ▪ feed extra ration along with mineral and vitamin supplements to withstand cold wave | <ul style="list-style-type: none"> ▪ Analysis of the present experience and remodeling of housing structure. |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ Selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. |
| Goat/Sheep | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Feed balance ration to withstand the cold wave prior to occurrence. | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to cold wave ▪ provide extra bedding materials ▪ feed extra ration along with mineral and vitamin supplements to withstand cold wave | <ul style="list-style-type: none"> ▪ Analysis of the present experience and remodeling of housing structure. |
| Health and disease | <ul style="list-style-type: none"> ▪ Ensure livestock insurance | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and |

| | | | |
|-------------------|--|--|---|
| management | <ul style="list-style-type: none"> ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <p>symptomatically prompt treatment accordingly.</p> <ul style="list-style-type: none"> ▪ Supplementary feeding of vitamin and mineral to improve general body health. | <p>symptomatically prompt treatment accordingly.</p> <ul style="list-style-type: none"> ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ Selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. |
| Snowfall | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. |
| Earthquake | NA | NA | NA |
| Landslides | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ immediate rescue operation ▪ Shifting of livestock to safe areas. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to landslides and remedies to be taken accordingly for future. |

^s based on forewarning wherever available

2.5.2 Poultry

| | Suggested contingency measures | | | Convergence/linkages with ongoing programs, if any |
|-------------------------------|--|--|--|--|
| | Before the event | During the event | After the event | |
| Drought | | | | |
| Shortage of feed ingredients | <ul style="list-style-type: none"> ▪ Awareness on maize, pea and oil seed cultivation for use of poultry feed ▪ Procurement of feed ingredients in bulk. ▪ Installation of feed mixing plant | <ul style="list-style-type: none"> ▪ Use of stored feed ▪ Use of feeds from the local resources ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Availing insurance for the crop loss. ▪ Availing subsidiary schemes from line deptt. | Schemes from Line Deptt./RKVY/ATMA |
| Drinking water | <ul style="list-style-type: none"> ▪ Construction of water harvesting structures. ▪ Harvesting rain water & water from natural source ▪ Developing watershed areas. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Provision of potable water ▪ Use of stored water from water harvesting structure. ▪ Fetching water from watershed areas and natural stream/river. ▪ Avail subsidy water supply through tankers from state or central Govt. | <ul style="list-style-type: none"> ▪ Submitting a memorandum to state or central Govt. regarding amount of water shortfall during drought and action to be initiated accordingly. ▪ Construction of permanent water harvesting structure with a planning to fulfill the water requirement during drought. | |
| Health and disease management | <ul style="list-style-type: none"> ▪ Regular deworming and vaccination against viral disease. ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. ▪ Proper ventilation system of Housing to reduce heat stress. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to reduce heat stress ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ selective culling of bird ▪ Submitting a memorandum to state or central Govt. regarding the loss of poultry due to Drought and remedies to be taken accordingly for future. | |
| Floods | | | | |

| | | | | |
|-------------------------------|--|--|--|----|
| Shortage of feed ingredients | <ul style="list-style-type: none"> ▪ Awareness on maize, pea and oil seed cultivation for use of poultry feed ▪ Procurement of feed ingredients in bulk and store in raised floor. ▪ Installation of feed mixing plant | <ul style="list-style-type: none"> ▪ Use of stored feed ▪ Use of feeds from the local resources ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Availing insurance for the crop loss. ▪ Availing subsidiary schemes from line deptt. | |
| Drinking water | <ul style="list-style-type: none"> ▪ Storage of safe drinking water in community tanks / water harvesting structures which is not prone to seepage of flood water. ▪ Installation of large sized sand filters with charcoal. ▪ Tying up with PHED Deptt. of neighboring district to supply water at needy time. ▪ Creating awareness amongst public how to conserve water and judiciously use in flood situation. | <ul style="list-style-type: none"> ▪ Chlorination of the drinking water and use of sand filter ▪ Supply of fresh drinking water from nearby district. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Cleaning of water storage tanks ▪ Relief for damaged tanks and community pipe line for reconstruction. | |
| Health and disease management | <ul style="list-style-type: none"> ▪ Regular deworming and vaccination against viral disease. ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. ▪ Proper ventilation system of Housing to reduce heat stress. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to reduce heat stress ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ selective culling of bird ▪ Submitting a memorandum to state or central Govt. regarding the loss of poultry due to Drought and remedies to be taken accordingly for future. | |
| Cyclone | | | | |
| Shortage of feed | NA | NA | NA | NA |

| | | | | |
|--------------------------------|--|---|--|----|
| ingredients | | | | |
| Drinking water | NA | NA | NA | NA |
| Health and disease management | NA | NA | NA | NA |
| Heat wave | | | | |
| Shelter/environment management | <ul style="list-style-type: none"> ▪ Advance early warning system through Agromet advisories for preparedness to combat the situation. ▪ Good shelter with well ventilation and bedding materials ▪ Construction of shelters in wind shed areas. ▪ Increase the concentrate feed amount and reduce the roughage diet. ▪ Adlib provision of potable water | <ul style="list-style-type: none"> ▪ Confine the animal in protected shelter ▪ prevent them direct expose to heat wave ▪ reduce upto 20% of the ration ▪ provide nutretical ▪ Adlib provision of potable water ▪ Avoid movement of animal ▪ Misting of water during the extreme heat to the animal | <ul style="list-style-type: none"> ▪ Adlib provision of potable water ▪ Analysis of the present experience and remodeling of housing structure. ▪ provide nutretical | |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ selective culling of injured animal | <ul style="list-style-type: none"> ▪ Immediate attention to the ailing animals. ▪ selective culling of injured animal ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Sanitization of the shed and surrounding areas. ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to flood and remedies to be taken accordingly for future. | |
| Cold wave | | | | |
| Shelter/environment | <ul style="list-style-type: none"> ▪ Good shelter with well | <ul style="list-style-type: none"> ▪ Confine the bird in protected | | |

| | | | | |
|-------------------------------|---|---|--|----|
| management | <p>ventilation and bedding materials</p> <ul style="list-style-type: none"> ▪ Construction of shelters in wind shed areas. ▪ Feed balance ration to withstand the cold wave prior to occurrence. | <p>shelter</p> <ul style="list-style-type: none"> ▪ prove extra light to keep them warm ▪ prevent them direct expose to cold wave ▪ provide extra bedding materials ▪ feed extra ration along with mineral and vitamin supplements to withstand cold wave ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <p>Analysis of the present experience and remodeling of housing structure.</p> | |
| Health and disease management | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load and vaccination to protect viral disease ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to cold wave and remedies to be taken accordingly for future. | |
| Snowfall | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load and vaccination to protect against viral disease ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to snow | NA |

| | | | | |
|----------------------------|---|---|---|----|
| | <ul style="list-style-type: none"> ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | | fall and remedies to be taken accordingly for future. | |
| Earthquake, Landslides etc | <ul style="list-style-type: none"> ▪ Ensure livestock insurance ▪ Deworming to reduce worm load and vaccination to protect against viral disease ▪ Stocking of veterinary medicines, vitamin and mineral supplements. ▪ Training of paravets and identifying key man in each village to combat the situation if arise. ▪ Providing available communication and transportation facilities in every dispensary / clinic for consultations. | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Supplementary feeding of vitamin and mineral to improve general body health. ▪ immediate rescue operation ▪ Shifting of livestock to safe areas. ▪ Regular radio/TV telecast to follow the instruction of Do & Don'ts from experts | <ul style="list-style-type: none"> ▪ Mass awareness cum Health camp and symptomatically prompt treatment accordingly. ▪ Immediate attention to the ailing animals. ▪ Sanitization of the shed and surrounding areas. ▪ selective culling of animal ▪ Submitting a memorandum to state or central Govt. regarding the loss of animal due to landslides and remedies to be taken accordingly for future. | NA |

^a based on forewarning wherever available